

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Licchtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau




*Handwritten signature*

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/12, 15/62, 5/10, C07K 14/435, 14/72, C12Q 1/02, G01N 33/50, 33/52</b>		A1	(11) International Publication Number: <b>WO 97/20931</b>
			(43) International Publication Date: 12 June 1997 (12.06.97)
(21) International Application Number: PCT/US96/19516		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 December 1996 (06.12.96)		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 60/008,373 8 December 1995 (08.12.95) US			
(71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; National Institutes of Health, Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HTUN, Han [US/US]; 4937 Ten Mills Road, Columbia, MD 21044 (US). HAGER, Gordon, L. [US/US]; 4525 Cambria Avenue, Garrett Park, MD 20896-1321 (US).			
(74) Agents: SELBY, Elizabeth et al.; Needle & Rosenberg, Suite 1200, 127 Peachtree Street, N.E., Atlanta, GA 30303 (US).			

(54) Title: METHOD AND COMPOSITIONS FOR MONITORING DNA BINDING MOLECULES IN LIVING CELLS

Structure of the MMTV tandem array in Cell Line 3134



(57) Abstract

The present invention provides a method of screening for a compound that binds to a selected nucleic acid comprising contacting compound fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of the nucleic acid in an array such that the nucleic acid can be directly detected when bound by fluorescently labeled compound; and directly detecting the location of fluorescence within the cell, fluorescence aggregated at the site of the nucleic acid array indicating a compound that binds to the selected nucleic acid. In particular compounds such a transcription factor can be screened. Reagents for such method are provided including a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and a chimeric protein comprising a fluorescent protein fused to a steroid receptor.

## METHOD AND COMPOSITIONS FOR MONITORING DNA BINDING MOLECULES IN LIVING CELLS

### BACKGROUND OF THE INVENTION

5

#### Field of the Invention

The present invention relates to the field of binding of molecules such as transcription factors to regions of nucleic acids, steroid hormone usage, steroid receptors and their corresponding response elements. Reagents are provided to allow methods involving direct detection of binding of a molecule, determining response element targeting by activated steroid receptors, screening for steroid agonists and antagonists, and monitoring levels of steroid agonists and antagonists in biological samples.

15 

#### Background Art

Steroid receptors are hormone-dependent activators of gene expression. Steroid receptors mediate the action of steroid hormones (*e.g.*, glucocorticoids, estrogens, progestins, testosterone, mineralocorticoids and 1,25-dihydroxycholecalciferol) in human tissues. After activation with the cognate ligand, receptors bind to chromatin in the nucleus and modulate the activity of target cellular genes. The binding of receptors to these target sequences is a key step in steroid function. Currently, this interaction can only be detected by indirect methods, such as reporter assays that detect the result of transcriptional activation coupled with transfection methods that introduce DNA sequences with receptor binding sites.

25

It is generally accepted that the unliganded glucocorticoid receptor (GR) resides in the cytoplasm, and that hormone activation leads both to nuclear accumulation and gene activation. (Gasc, J. -M. & Baulieu, E. E. (1987) in *Steroid Hormone Receptors: Their Intracellular Localisation*, ed. Clark, C. R. (Ellis Horwood Ltd., Chichester, England), pp. 233-250; Beato, M. (1989) *Cell* **56**, 335-344; Carson-Jurica, M. A., Schrader, W. T. & O'Malley, B. W. (1990) *Endocr. Rev.* **11**, 201-220; Gronemeyer, H. (1993) in *Steroid Hormone Action*, ed. Parker, M. G. (Oxford University Press, New York), pp. 94-117; Tsai, M. J. & O'Malley, B. W. (1994) *Annu. Rev. Biochem.* **63**,

451-486; Akner, G., Wikstrom, A. C. & Gustafsson, J. A. (1995) *J. Steroid Biochem. Mol. Biol.* **52**, 1-16), and references therein. However, the mechanisms involved in nuclear translocation and targeting of steroid receptors to regulatory sites in chromatin have been poorly understood. It has previously been difficult to discriminate between  
5 the ability of a given receptor mutant, or a given receptor/ligand combination, to participate in the separate processes of receptor activation, nuclear translocation, sequence-specific binding, and promoter activation.

Proteins have previously been labeled with fluorescent tags to detect their  
10 localization and sometimes their conformational changes both *in vitro* and in intact cells. Such labeling is essential both for immunofluorescence and for fluorescence analog cytochemistry, in which the biochemistry and trafficking of proteins are monitored after microinjection into living cells (Wang, Y.L. & Taylor, D.L., eds. (1989) *Methods Cell Biol.* **29**). Traditionally, fluorescence labeling is done by purifying proteins and then  
15 covalently conjugating them to reactive derivatives of organic fluorophores. The stoichiometry and locations of dye attachment are often difficult to control, and careful repurification of the proteins is usually necessary. If the proteins are to be used inside living cells, a final challenging step is to get them across the plasma membrane via micropipet techniques or various methods of reversible permeabilization. Furthermore,  
20 in previous hormone studies broken cell preparations or antibody tags in fixed cell preparations were used, both techniques that cause enormous disruption of cell structures.

The green fluorescent protein (GFP) from the jellyfish *Aequorea victoria* is a  
25 molecule whose natural function seems to be to convert the blue chemiluminescence of the  $\text{Ca}^{2+}$ -sensitive photoprotein aequorin into green emission (Ward, W.W. (1979) in *Photochemical and Photobiological Reviews*, ed. Smith, K.C. (Plenum, New York), 4:1-57). GFP's absorption bands in the blue (maximally at a wave length of 395nm with weaker absorbance at 470nm) and emission peak in the green (at 509nm) do not arise  
30 from a distinct cofactor but rather from an internal *p*-hydroxybenzylideneimidazolidinone chromophore generated by cyclization and

- oxidation of a serine-tyrosine-glycine sequence at residues 56-67 (Cody, C.W., Prasher, D.C., Westler, W.M., Prendergast, F.G. & Ward, W.W. (1993) *Biochemistry* **32**, 1212-1218). The gene for GFP was cloned (Prasher, D.C., Eckenrode, V.K., Ward, W.W., Prendergast, F.G. & Cormier, M.J. (1992) *Gene* **111**, 229-233), and the encoded
- 5 protein consists of 238 amino acid residues (molecular weight 27kD). Heterologous expression of the gene has been done in *Escherichia coli* (Heim, R., Prasher, D.C. and Tsien, R.Y. (1994) *Proc. Natl. Acad. Sci. U.S.A.* **91**, 12501-12504); Inouye, S. & Tsuji, F.I. (1994) *FEBS Lett.* **341**, 277-280), *Caenorhabditis elegans* (Chalfie, M., Tu, Y., Euskirchen, G., Ward, W.W. & Prasher, D.C. (1994) *Science* **263**, 802-805), and
- 10 *Drosophila melanogaster* (Yeh, E., Gustafson, K. & Boulianne, G.L. (1995) *Proc. Natl. Acad. Sci. U.S.A.* **92**, 7035-7040; Tannahill, D., Bray, S. & Harris, W.A. (1995) *Dev. Biol.* **168**, 694-697 and plants (Hu, W. & Cheng, C.L. (1995) *FEBS Lett.* **369**, 331-334; Baulcombe, D.C., Chapman, S. & Santa Cruz, S. (1995) *Plant J.* **7**, 1045-1053). Recently, chimeric genes encoding N- and C-terminal fusions of the *Drosophila*
- 15 exuperantia (exu) gene product, Exu (Wang, S. and Hazelrigg, T. (1994) *Nature* **369**, 400-403), actin Act88F gene (Barthmaier, P. and Fyrberg, E. (1995) *Dev. Biol.* **169**, 770-774), and a nuclear localization signal (Davis, I., Girdham, C.H. & O'Farrell, P.H. (1995) *Dev. Biol.* **170**, 726-729); of the yeast microtubule and spindle pole associated dis1 gene product (Nabeshima, K., Kurooka, H., Takeuchi, M., Kinoshita, K.,
- 20 Nakaseko, Y., & Yanagida, M. (1995) *Genes Dev.* **9**, 1572-1585) and an RNA binding protein Npl3p (Corbett, A.H., Koepp, D.M., Schlenstedt, G., Lee, M.S. Hopper, A.K. & Silver, P.A. (1995) *J. Cell Biol.* **130**, 1017-1026); and of a mammalian ion channel protein, NMDAR1 (Marshall, J., Molloy, R., Moss, G.W., Howe, J.R. & Hughes, T.E. (1995) *Neuron* **14**, 211-215), microtubule-associated protein, MAP4 (Olson, K.R.,
- 25 McIntosh, J.R. & Olmsted, J.B. (1995) *J. Cell Biol.* **130**, 639-650), and a secretory protein, chromogranin B (Kaether, C. & Gerdes, H.H. (1995) *FEBS Lett.* **369**, 267-271) have been constructed fused to GFP. However, none of these chimeric proteins have been to transcription factors or co-factors and no suggestions have been made as to the usefulness of such a fusion to study physiologically relevant interaction on an amplified

DNA target. Furthermore, none of these reports indicated a successful use of GFP in mammalian cells.

Many human diseases result from aberrant steroid function, and many disease  
5 states, *i.e.*, inflammation, are treated with glucocorticoid and other steroid derivatives. A large number of drugs have been developed whose function is based on the ability to interact with and activate steroid receptors. The identification and characterization of these compounds is a laborious, time-consuming and expensive process involving years of work. Even with a large investment of resources, the true behavior of these  
10 compounds in living cells is not understood.

The present invention allows observation for the first time of *in vivo* target sites within a higher eukaryotic nucleus for trans-regulatory molecules, such as transcription factors, *e.g.*, glucocorticoid receptor (GR). The visualization of physiologically  
15 relevant *in vivo* target sites for any transcription factor to date has not previously been accomplished. The present invention provides a powerful method for identification of any single target site in a higher eukaryotic genome, comprising roughly 60,000-80,000 genes (Bird, A.P. (1995) *Trends Genet.* 11:94-100), using a singly fluorescently-labelled regulatory factor, which has not been considered previously. Discriminating direct  
20 versus indirect interaction between a regulatory molecule and its putative regulatory site is critical for the development of highly specific drugs directed against trans-regulatory factors. Traditionally, the methodology for showing potentially direct interactions involves nuclease or chemical protection experiments and transient co-transfection experiments of the putative regulator and its regulated site. While this approach  
25 indicates potential direct interaction, it does not necessarily imply direct interaction. Alternatively, the approach of making compensatory mutations between the regulatory sequences as well as the DNA binding specificity has been used in an attempt to demonstrate direct regulatory interaction (Schier, A.F. and Gehring, W.J. (1992) *Nature* 356:804-807), an extension of the principles of second site suppression in genetics to  
30 molecular biology. However, such an approach makes enormous assumptions of our understanding of sequence-specific recognition by sequence-specific DNA binding

proteins *in vivo*, which certainly would not be valid for many systems, since many profound developmental events are governed by exquisite interactions to fine tune the system regarding, for example, concentration gradients of trans-regulatory factors. The present invention allows a simple and straight-forward manner in which direct  
5 interaction between a sequence-specific DNA binding protein or its co-factor and its putative regulatory site in the *in vivo* genomic context can be addressed. With this simple inventive methodology, novel classes of drugs directed not only against members of the steroid-ligand-dependent transcription factors but to new classes of drugs that target other transcription factors or their co-factors can be screened.

10

Additionally, the present invention provides the first opportunity to observe and monitor gene targeting specifically of steroid receptors in living cells wherein binding of the steroid receptor to its response element target can be observed distinctly from translocation of steroid receptor. The invention therefore provides for many relevant  
15 analyses, such as real-time determination of steroid activity in subjects as well as screening of compounds for response element binding/targeting capabilities as distinct from translocation capabilities. Such methods have implications in many diseases associated with steroid hormones, such as endocrine disorders, rheumatic disorders, collagen diseases, dermatological diseases, allergic states, ophthalmic diseases,  
20 respiratory disease, hematologic disorders, neoplastic disease, edematous states, gastrointestinal diseases and neurological conditions, and in other uses such as prevention of rejection of transplanted tissues.

## SUMMARY OF THE INVENTION

The present invention provides a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly  
5 detected when bound by fluorescently labeled steroid receptor.

The present invention further provides a chimeric protein comprising a fluorescent protein fused to a transcription factor. The present invention also provides a chimeric protein comprising a fluorescent protein fused to a steroid receptor.

10

The instant invention provides an isolated nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a transcription factor and an isolated nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a steroid receptor.

15

The instant invention also provides a cell containing a nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a transcription factor and a cell containing a nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a steroid receptor.

20

The instant invention provides a method of screening for a compound that binds to a selected nucleic acid comprising:

- a. contacting compound fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of the nucleic acid in an array such that  
25 the nucleic acid can be directly detected when bound by fluorescently labeled compound; and
- b. directly detecting the location of fluorescence within the cell, fluorescence aggregated at the site of the nucleic acid array indicating a compound that binds to the selected nucleic acid.

30



The present invention also provides a method of characterizing a ligand's effect on cellular localization of a compound to which the ligand binds in a cell comprising:

- a. contacting the ligand with a cell having the compound fluorescently labeled by a fluorescent protein and
  - 5        b. directly detecting the location of fluorescence within the cell,
- the location of fluorescence in the cell indicating the localization effect of the ligand on the compound.

Additionally provided is a method of determining a binding site for a DNA-  
10 binding protein comprising:

- a. contacting the DNA-binding protein fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of a nucleic acid having a putative binding site in an array such that the putative binding site can be directly visualized when bound by the fluorescently  
15        labeled DNA-binding protein, and
  - b. directly detecting the location of fluorescence within the cell,
- the presence of fluorescence aggregated at the putative binding site indicating a binding site to which the DNA-binding protein binds.

20        The present invention also provides a method of screening for a ligand that activates gene targeting of a steroid receptor in the nucleus of a mammalian cell comprising:

- a. contacting the ligand with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element  
25        can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
- b. directly detecting the location of fluorescence within the cell,

fluorescence aggregated at the site of the steroid receptor response element array in the nucleus indicating a ligand that activates the gene targeting of a steroid receptor in the nucleus of a mammalian cell.

- 5       The present invention provides a method of screening for a ligand that activates the translocation of a steroid receptor to the nucleus in a mammalian cell comprising:
- a.       contacting the ligand with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid  
10       receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
  - b.       directly detecting the location of fluorescence within the cell,  
the location of fluorescence aggregated in the nucleus indicating a ligand that activates  
15       the translocation of a steroid receptor to the nucleus in a mammalian cell.

The instant invention provides a method of detecting in a biological sample the presence of an agonist of a steroid receptor comprising:

- a.       contacting the sample with a mammalian cell having a plurality of steroid  
20       receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
- b.       directly detecting the location of fluorescence within the cell,  
25       the location of fluorescence aggregated at the site of the steroid receptor response element array in the nucleus indicating the presence of an agonist of the steroid receptor in the sample.

- 30       The present invention also provides a method of detecting in a biological sample the presence of an antagonist of a steroid receptor comprising:

- a. contacting the sample and an agonist of the steroid receptor with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
- 5 b. directly detecting the location of fluorescence within the cell,
- the absence of fluorescence substantially aggregated at the site of the steroid receptor response element array in the nucleus indicating the presence of an antagonist of the
- 10 steroid receptor in the sample.

The present invention provides a method of monitoring the level of an agonist of a steroid receptor in a subject comprising:

- a. periodically obtaining a biological sample from the subject,
- 15 b. contacting the sample with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid
- 20 receptor; and
- c. directly detecting the location of fluorescence within the cell,
- a decrease in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating a decrease in level of the steroid agonist of the steroid receptor in the sample
- 25 and an increase in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid agonist of the steroid receptor in the sample.

The instant invention provides a method of monitoring the balance between

30 levels of an agonist of a steroid receptor and an antagonist of the steroid receptor in a subject comprising:

- a. periodically obtaining a biological sample from the subject,
  - b. contacting the sample with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
  - c. directly detecting the location of fluorescence within the cell,
- an increase in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid agonist relative to level of the steroid antagonist in the sample, and a decrease in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid antagonist of the steroid receptor relative to level of the steroid agonist in the sample.

The instant invention also provides a method of determining an effective dosage of a steroid receptor agonist in a subject comprising:

- a. transferring into a set of cells from the patient a nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a steroid receptor;
- b. contacting the cells in the set with one of a selected range of dosages of the steroid agonist; and
- c. directly detecting location of fluorescence in the set of cells, a dosage capable of locating fluorescence substantially in the nucleus indicating an effective dosage of steroid receptor agonist.

The present invention provides a method of determining an effective dosage of a steroid receptor agonist to maintain steroid receptor activation for a selected period of time in a subject comprising:

- a. administering to the subject a dosage of the steroid receptor agonist,

- b. periodically obtaining a biological sample from the subject,
  - c. contacting the sample with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
  - d. directly detecting the location of fluorescence within the cell,
- a dosage that maintains the location of fluorescence at the site of the steroid receptor response element array in the nucleus for the selected period of time indicating an effective dosage.

The present invention also provides a method of determining an effective dosage of a steroid receptor antagonist to abrogate agonist activity for a selected period of time in a subject comprising:

- a. administering to the subject a dosage of the steroid receptor agonist,
  - b. periodically obtaining a biological sample from the subject;
  - c. contacting the sample with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and
  - d. directly detecting the location of fluorescence within the cell,
- a dosage that prevents the location of fluorescence at the site of the steroid receptor response element array in the nucleus for the selected period of time indicating an effective dosage.

The present invention also provides a method of detecting a defect in a response pathway of a steroid receptor in a subject comprising transferring into a cell from the subject a nucleic acid functionally encoding a chimeric protein comprising a fluorescent

protein fused to the steroid receptor and detecting the location of fluorescence within the cell as compared to the location of fluorescence within a normal, control cell transfected with the nucleic acid, a difference in location of fluorescence within the cell of the subject as compared to location of fluorescence within the normal, control cell  
5 indicating a defect in the response pathway of the steroid receptor.

The instant invention provides a method of determining whether a defect in a response pathway of a steroid receptor in a subject is in translocation of the steroid receptor to a cell nucleus, comprising transferring into a cell from the subject having the  
10 defect a nucleic acid functionally encoding a chimeric protein comprising a fluorescent protein fused to the steroid receptor and detecting the location of fluorescence within the cell, the location of fluorescence substantially in the cytoplasm of the cell indicating the defect is in translocation of the steroid receptor to the nucleus.

## 15 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram of the integrated, tandem array of bovine papilloma virus (BPV) 69% transforming fragment, mouse mammary tumor virus (MMTV) long terminal repeat (LTR) containing glucocorticoid receptor (GR) binding sites, and  
20 Harvey murine sarcoma virus v-ras sequences (Ha-v-ras).

Figure 2 shows, in panels 11-17, the appearance of GFP-GR when the fluorescent tag is excited by 489 nm laser light, and the 511 nm emission examined by confocal microscopy, using a standard fluorescein filter set. Individual sections are depicted for a representative 3134 cell nucleus. As one focuses on 0.18 micrometer  
25 sections throughout the nucleus, a continuous fiber of intense light emission is detected over 5-7 sections, which corresponds to GFP-GR binding to the continuous BVP/MMTV-LTR/ras array. The array is seen in sections 12-16. Below the panels is provided a schematic presentation of the appearance of GFP-GR when the fluorescent protein is excited.

30 Figure 3 shows the construction of GFP-GR. (A) Plasmid pCI-nGFP-C656G contains the green fluorescent protein fused to the C656G mutant glucocorticoid

receptor. (B) Dexamethasone (dex) dependent stimulation of MMTV-pLTRLuc is shown for GFP-GR transfected cells. Solid bars represent the IL2R<sup>+</sup> selected population activated with 1 nM dex, and the open bar depicts activation of the endogenous receptor with 100 nM dex. (C) Ligand specificity is presented for activation of endogenous  
5 MMTV-LTR-CAT sequences present in the 1471.1 cells. Cells were treated for 4 hrs with the indicated ligand, then harvested and levels of CAT activity determined.

Figure 4 shows nuclear localization of GFP-GR in MCF7 cells when the cells are treated with (A) dexamethasone or (B) RU486. The fluorescent tag is excited by 489 nm laser light, and the 511 nm emission examined by confocal microscopy, using a  
10 standard fluorescein filter set.

Figure 5 shows nuclear localization of GFP-ER in (A) MCF7 cells and (B) MDA-MB-231 cells when the fluorescent tag is excited by 489 nm laser light, and the 511 nm emission examined by confocal microscopy, using a standard fluorescein filter set. MCF7 cells are reported to be estrogen receptor positive and hormone dependent.  
15 MDA-MB231 cells are reported to be estrogen receptor negative and hormone independent.

### DETAILED DESCRIPTION OF THE INVENTION

20 The present invention provides cell lines such as the murine cell line 3134, that contains a set of mouse mammary tumor virus (MMTV) Harvey murine sarcoma virus (HaMuSV) v-ras sequences organized in a head-to-tail tandem array of approximately 200 copies. Each MMTV promoter sequence in this array contains 4 glucocorticoid receptor (GR) binding sites; the complete array thus contains nearly 1000 GR binding  
25 sites. This cell is used to visualize directly the interaction between GR and its binding site in chromatin in living cells. This is accomplished with a fluorescent labeled copy of the GR. This array thus provides the unique opportunity to visualize direct receptor/target interactions. This structure, for which there is no precedent, provides for the first time a reagent to observe the interaction of steroid receptors with their  
30 response elements in living cells, and to characterize the effectiveness of medically important steroid ligands in activating gene expression in mammalian cells.

In the present application, we have shown the usefulness of GFP in monitoring the activity of a steroid hormone receptor. The ability to directly observe living cells has allowed us to follow in real time the process of cytoplasm to nuclear translocation, and has revealed for the first time differences in GR intranuclear accumulation pattern  
5 dependent on the type of activating ligand. Furthermore, the patterns of GR accumulation are remarkably similar between adjacent cells, suggesting an order in the organization of the interphase nucleus. This ability to observe living cells has also revealed subcellular localization of partially activated estrogen receptor (ER). Thus, the use of GFP has revealed new details about steroid localization and organization of the  
10 eukaryotic nucleus.

As used in the specification and in the claims, "a" can mean one or more, depending upon the context in which it is used.

15 The present invention provides reagents and methods for detecting, by direct visual evaluation, the binding of fluorescently labelled compounds to a nucleic acid. This direct detection can be accomplished by the use of a cell line having a sufficient number of copies of the binding region of the nucleic acid in an array, such as tandem repeats, that allows detection of the array when a fluorescently labeled compound is  
20 bound to the binding region by direct detection of the fluorescence localized at the site in the cell nucleus of the nucleic acid array. For example, after binding, cells can immediately, without further treatment of the cells, be placed under a fluorescent microscope and fluorescence directly visualized. Thus compounds can be rapidly analyzed for binding capability in a manner that clearly depicts the binding. Importantly,  
25 the binding detected in this assay is biologically relevant. The detected event represents receptor mobilization to a correct genetic target in the living cell. Thus, interactions revealed with this assay are much more reliable as measures of biologically germane receptor activity. Additionally, analyses of levels of compounds or defects in pathways involving the binding of such compounds to their nucleic acid binding site in specific  
30 subjects can be performed, as further described below.



Any selected nucleic acids and binding sites, and compounds that may bind thereto, either directly or indirectly, can be analyzed by this method, in a desired cell, as elaborated below. For example, binding of any transcription factor to its activation site on a nucleic acid can be directly determined along with the proteins associated with the factor (e.g., co-activator, co-repressor, adapter, or molecules in a similar category). Additionally, for example, binding of any selected steroid receptor can be directly detected and analyzed for both translocation to the nucleus and binding to the corresponding response element(s) in the nucleus. Thus, compounds typically can be proteins, polypeptides and peptides; however, other compounds can include, for example, Peptide Nucleic Acids (PNAs), antisense nucleic acids and organic molecules (e.g., dexamethasone). Importantly, for any compound, the step of binding to the binding site can visually be distinguished from the step of translocation to the nucleus, or alternatively, exit from the nucleus.

In general, the present invention utilizes fluorescent labeling of the compound by a fluorescent protein, as fluorescent protein is herein described, adding the labeled compound to cells, and directly detecting the location and/or aggregation of fluorescence in the cells. For detection of the translocation of the labeled compound to the nucleus, any cell can be utilized, since the resulting location of fluorescence can be visualized as either in the cytoplasm or in the nucleus. Additionally, for such detection events, cells having increased copy number of the binding site, in any array, can be used. For detection of binding to the target nucleic acid site, the present invention provides cells having a plurality of nucleic acid binding sites in an array such that the nucleic acid binding site can be directly detected when bound by binding compound, such as a ligand, transcription factor, etc., fluorescently labeled by a fluorescent protein, as described herein.

As used herein, direct detection means detection of the fluorescence emitted from the site in the cells when excited by light, ultraviolet or visible, without the need for any additional chemical reactions or treatment of the cells. The fluorescence is directly detected by any device capable of detecting fluorescence, such as a fluorescent

microscope, as visualized by the eye of the operator of the microscope at the time or as recorded from the microscope such as by photography of the field of view or through the use of photosensitive detectors. A fluorescent microscope, such as a confocal laser scanning microscope or an epifluorescent microscope, can be used, as is known in the art. There is no requirement that cells be, *e.g.*, fixed or stained or contacted by any additional reagents, in order to detect the binding. Thus living cells can be assayed, and results obtained, immediately after binding. Therefore, for example, subjects can be advised immediately of results of analyses as described below. Furthermore, it is anticipated that screening of both nuclear localization of fluorescence (translocation) and focal localization of fluorescence on a target array will be adapted to high volume computerized image analysis. That is, the analysis of large numbers of samples will be automated for either the repetitive examination of clinical samples or the large-scale screening of compounds in the research environment.

The cell can be derived from any desired mammal, such as, for example, human, monkey, mouse, hamster and rat. The nucleic acid can be amplified in an appropriate array by any of several means, as known to those skilled in the art. Generally, a selected nucleic acid binding site or collection of sites, for example as found within the context of a transcriptional regulatory region, *i.e.*, promoters, enhancers, silencers, etc., can be amplified in an array detectable by the present means, for example, by gene amplification of the nucleic acid binding site (*e.g.*, the steroid receptor response element or the transcription factor binding site) by linking it to a gene readily amplified in a tandem array, for example, dihydrofolate reductase, or by multimerization of the nucleic acid binding site or sites by synthetic DNA synthesis and/or enzymatic synthesis, for example, through the use of ligase and polymerases, and introducing the amplified element into selected mammalian cells. Such methods are further elaborated in the examples provided below.

The cells used herein have an array of the nucleic acid having binding sites under analysis such that the nucleic acid can be directly detected when bound by a fluorescently labeled compound. Such array as contemplated herein has sufficient copies

of the nucleic acid and in such an arrangement that the fluorescently labeled binding compound, when bound to the site can be directly detected and readily identified. Therefore, the array includes arrangement of the copies in sufficiently close physical proximity along a chromosome, either present endogenously or artificially introduced or  
5 induced, or in extrachromosomally replicating episomes, to allow localization of fluorescence at a discrete, detectable site in the nucleus, as seen under standard magnification for cells and nuclei. Such array as contemplated herein allows detection in the context of chromatin, as exists in the interphase nucleus. An example of such an array is a series of direct tandem repeats of the nucleic acid. An example of a tandem  
10 array of direct repeating units is depicted in FIG. 1.

By a "plurality" of any herein described nucleic acid having a binding site is meant that the number of copies of the nucleic acid having the binding site (e.g., the steroid receptor response element or the transcription factor binding site) is greater than  
15 one. Preferably, the cells have more than about five copies, more preferably more than about ten copies, more preferably more than about twenty, and more preferably more than about forty copies. For example, cell line 3134, described herein, has about two hundred copies of the MMTV LTR-Ha-v-ras- gene, each of which has four copies of the binding site for glucocorticoid receptor and with each site accommodating two  
20 glucocorticoid receptor molecules. Any number which allows detection of the site upon binding of the fluorescently labeled binding compound is contemplated. Thus, an example of a cell of the present invention is a cell of the cell line 3134 deposited with American Type Culture Collection as accession number CRL-11998 (ATTC).

25 Specifically, the present invention provides a cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor. A response element, as used herein, includes any nucleic acid to which a steroid receptor directly binds, but also includes the steroid receptor associated, either directly or indirectly, factors that are  
30 recruited to the vicinity of the element (e.g., nuclear factor 1 (NF1), octomer transcription factor 1 (OTF1), steroid receptor coactivator 1 (SRC1), etc.). Steroid

receptors, and corresponding response elements to which they bind, can include any steroid receptor, for example, glucocorticoid receptor, estrogen receptor (ER), progesterone receptor, androgen receptor, mineralocorticoid receptor, vitamin D receptor. Examples of steroid receptor response elements include those contained in the

5 mouse mammary tumor virus (MMTV) long terminal repeat (LTR) (which has binding sites for glucocorticoid receptor, mineralocorticoid receptor, progesterone receptor, and androgen receptor), and those contained in vitellogenin and osteocalcin genes (which have binding sites for estrogen and vitamin D receptors). Thus, response elements in such array in a cell can include other transcriptional regulatory elements contained

10 within the mouse mammary tumor virus long terminal repeat and bovine papilloma virus 69% transforming DNA. Many steroid receptors and steroid response elements, as exemplified above, are known to the skilled artisan; however, any steroid receptor and its response element is contemplated herein. In addition to steroid receptors, there are other ligand-dependent receptors (such as thyroid hormone receptor, retinoic acid

15 receptor, retinoid X receptor, TCDD (dioxin) receptor, fatty acid activatable receptors, etc.) and stimulus-dependent receptors (such as peroxisome proliferator activated receptor, growth factor-dependent receptors (e.g., epidermal growth factor, nerve growth factor, etc.)), and factors (such as CREB, NFAT, NFkB/IkB, etc.), and other receptors whose ligand remains to be defined (such as mammalian homologs of the

20 *Drosophila* tailless, knirps, sevenup, FTZFI genes, etc.). Many of these receptors or factors can be found listed in the book [Parker, M.G. (1993) *Steroid Hormone Action* (Oxford University Press, New York, pp. 210)], in a recent review article [Tsai, M.J. & O'Malley, B.W. (1994) *Annu. Rev. Biochem.* 63, 451-486], and in the GenBank database, which will contain additional receptors as well as the complete nucleotide

25 sequences of the genes and cDNAs. In addition, the cell line offers a number of tandemly repeated regulatory sites for sequence-specific transcription factors (such as activating protein 2 (AP2), OTF1, NF1/CTF, etc.) as well as general transcription factors (such as TFIID, initiator protein, etc.). The steroid receptor response elements (or other transcriptional regulatory elements) used in the present invention in arrays

30 detectable as described herein can be integrated into the genome of the cell, maintained

in the cell on artificial mammalian chromosomes (Monaco, A.P. and Larin, Z. (1994) *Trends Biotechnol.* **12**, 280-286) or can be carried on episomal elements.

The cell can further comprise a nucleic acid encoding a chimeric protein wherein  
5 a fluorescent protein is fused to the steroid receptor. Steroid receptors as described  
above are contemplated for use in a chimeric protein. Typically, the steroid receptor in  
the chimeric protein encoded by the nucleic acid in a selected cell is one that binds to the  
response element in the selected cell in the herein described array. Throughout this  
application by "a fluorescent protein" is meant a protein that fluoresces in cells without  
10 adding exogenous cofactors. That is, it is a protein that can be expressed in cells and  
detected in these cells simply by exciting the protein with light and visualizing the  
resultant fluorescence. An example of such a fluorescent protein is the green fluorescent  
protein (GFP) originally isolated from the jellyfish *Aequorea victoria*. Another example  
of a fluorescent protein as defined herein is the green fluorescent protein originally  
15 isolated from *Renilla reniformis*, which demonstrated a single absorption peak at 498 nm  
and an emission peak at 509 nm. (Cubitt, *et al.* (1995) *TIBS* **20**: 448-455). By  
fluorescent protein is also contemplated that modifications may be made to a fluorescent  
protein, as long as the resulting protein fluoresces when expressed in cells.  
Modifications can be developed based upon the chemistry of chromophore formation.  
20 (Cubitt, *et al.* *TIBS* **20**: 448-455). Generally, however, one may prefer to leave the  
glycine which participates in forming the chromophore, in part by cyclization of a Ser-  
Tyr-Gly moiety (Gly 67 in *Aequorea* GFP in the Ser65-Tyr66-Gly67 moiety), intact. An  
example of a useful substitution that modifies the absorption spectra is the substitution  
in *Aequorea* green fluorescent protein for serine at amino acid 65 by, for example,  
25 threonine, cysteine, leucine, valine, or alanine, that allows the excitation of the  
chromophore at a lower energy (longer wavelength) than the naturally occurring protein  
thereby greatly decreasing the destruction of the chromophore as occurs when it is  
excited at a higher energy. Such mutation at amino acid 65 also increases brightness and  
rate of oxidation as compared to wild-type *Aequorea* GFP when each is excited at its  
30 longest wavelength peak. In addition, other spectral variants of GFP, such as improved  
blue variants of GFP have been developed (e.g., pCI-nGL2-C656G; pCI-nGL3-C656G;

pCI-nGL4-C656G; pCI-nGL5-C656G; pCI-nGL7-C656G; pCI-nGL9-C656G; pCI-nGL10-C656G; pCI-nGL11-C656G). Variants emitting longer wavelengths (*e.g.*, red variants) can also be developed by introducing other mutations into the GFP DNA. Additionally, the codon usage of any GFP- coding sequence can be modified to human codons, according to known methods. For example, pGreenLantern-1 (LifeTechnologies, Inc., Gaithersburg, MD, catalog number 10642-015) is a commercially available S65T variant GFP cDNA with mammalian codon usage. Other commercially available humanized GFP-cDNAs are: pEGFP-N (catalog numbers: 6086-1; 6085-1; 6081-1 from Clontech), pEGFP-C (catalog numbers: 6084-1; 6083-1; 6082-1 from Clontech), and pHGFP-S65T (catalog number 6088-1 from Clontech). Additional useful modifications of any fluorescent protein can include other modifications that speed up the rate of the oxidation step of chromophore formation, that increase brightness at longer wavelengths, and that reduce photoisomerization and/or photobleaching. Furthermore, in general it is preferable that GFP not be truncated by more than about one amino acid from the amino terminus and about 10-15 amino acids from the carboxyl terminus. Detection of additional fluorescent proteins can readily be performed by standard approaches such as searching for proteins having some homology to GFP in nucleic acid libraries from organisms that demonstrate fluorescence by nucleic acid hybridization and by searching for homologous nucleic acids and proteins in other organisms in databanks of nucleic acid and protein sequences and testing the encoded proteins for fluorescence. It is possible that forced protein evolution of the currently existing GFP can be achieved by randomizing the entire GFP coding region so as to make every single possible change at every single amino acid coding region as well as pairs and further combination of changes. Desirable changes yielding better chromophore or different excitation/emission spectra can be characterized by fluorescence spectroscopy or flow cytometry upon translation of the coding sequences into proteins. An example of one general approach would be to take advantage of the phage display system for expression of the chromophore on the surface of a bacteriophage using a modification of what is currently being done for antibodies (*e.g.* Pharmacia Biotech, Inc.'s Recombinant Phage Antibody System). Another general approach would be to adapt a protocol similar to that used to select novel enzymatic

activities displayed by RNAs (Bartel, D.P. and Szostak, J.W. (1993) *Science* **261**, 1411-1418) for selection of novel fluorescent proteins.

Therefore, also provided herein are chimeric proteins comprising a fluorescent  
5 protein fused to a transcription factor, and nucleic acids encoding such proteins. One  
example of a transcription factor that can be used herein, described in further detail  
above, is a steroid receptor; however, numerous other transcription factors can be  
utilized. For example, basal transcription factors (e.g. TFIID, etc.), and sequence  
specific DNA binding transcription factors (e.g., AP1, AP2, SP1, NF1, etc). Additional  
10 transcription factors are listed in, for example, computer databases such as that  
maintained by the National Center for Biotechnology Information (NCBI, Bethesda,  
MD) accessible through the BLAST program (see item 19 (TFD) for transcription  
factors; item 20 for eukaryotic promoter sequences). Additionally, as used in the claims,  
"transcription factors" include transcription adaptor molecules or cofactors, which  
15 localization within the cell can be monitored also by this method. Transcription adaptor  
molecules or cofactors are those molecules that interact with transcription factors to  
effect their function (*i.e.*, their activation or repression functions). For example, SRC1,  
steroid receptor coactivator 1, is a cofactor of steroid receptors.

20 The chimeric protein can include a linking peptide sequence between the  
fluorescent protein and the steroid receptor. For example, a sequence of the amino  
acids glycine and alanine, or a sequence of alanine alone can be used; however, any  
sequence of amino acids and any length can be used that does not interfere with the  
binding of the steroid receptor to its response element and that does not prevent  
25 fluorescence of the fluorescent protein. Typically, a linker peptide will range from two  
to about ten amino acids but maybe shorter or longer. Of course, certain linker peptides  
maybe preferred over others, e.g., the presence of four basic amino acids in a string of  
six might suffice as a nuclear localization signal so as to mislocate the uninduced state of  
the factor. A linker peptide can be used to separate the fluorescent protein structurally  
30 from the response element and can function to allow the fluorescent protein  
independently of the remaining portion of the chimeric protein. An example of a

chimeric protein, which has GFP fused via five glycine-alanine repeating units to the N-terminus of the glucocorticoid receptor, is provided herein as SEQ ID NO: 2. Typically, a fluorescent protein can be fused to either the C-terminus or the N-terminus of the transcription factor; however, the preferable construction for a specific transcription  
5 factor can readily be determined. Linker peptides can readily be introduced between the two proteins in the chimeric protein by producing an nucleic acid that encodes the chimeric protein having the linker sequence between the fluorescent protein and transcription factor components.

10 Modifications to the fluorescent protein portion of the chimeric protein and/or the transcription factor, e.g., steroid receptor can be made. For example, a green fluorescent protein can be modified as described above. The transcription factor, for example, can be modified to increase or decrease its affinity for its binding site or to determine if a selected modification affects its binding affinity. In the case of steroid- or  
15 ligand-dependent transcription factor, the region involved in steroid- or ligand-binding can be altered to either increase or decrease the affinity to the steroid or ligand or alter the specificity of the ligand. Furthermore, other functions of the factor, such as transactivation potential, maybe modified. An example of such a modification is found in the chimeric protein having the amino acid sequence set forth in SEQ ID NO: 2,  
20 wherein the transcription factor is glucocorticoid receptor having a substitution of serine for cysteine at amino acid 656 that has a higher binding affinity for its ligand than the protein having the naturally occurring amino acid sequence. This substitution also increases the transactivation potential of the receptor, resulting in "superactivation." This cysteine 656 mutation can be utilized, for example in rat, human and mouse  
25 glucocorticoid receptor. For example, steroid receptors, or any transcription factor, can be modified in their steroid binding domains to increase affinity for steroid, thus allowing one to increase use of exogenous receptor over endogenous receptor in a cell.

The transcription factor of the chimeric protein can be derived from any selected  
30 mammal. Additionally, chimeric proteins utilizing a transcription factor from one mammal can often be used in a cell from another mammal. For example, the



glucocorticoid receptor amino acid sequence is highly conserved, particularly in the binding region among rat, human and mouse, and, for example, the rat glucocorticoid receptor binds with high affinity to the human glucocorticoid response element.

5           Also provided in the present invention is nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to a transcription factor. The nucleic acid encoding the chimeric protein can be any nucleic acid that functionally encodes the chimeric protein. For example, to functionally encode, *i.e.*, allow the nucleic acid to be expressed, the nucleic acid can include, for example, expression control sequences, such  
10 as an origin of replication, a promoter, an enhancer, and necessary information processing sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from metallothionine genes, actin genes, immunoglobulin genes, CMV, SV40, adenovirus, bovine papilloma virus, etc. A nucleic acid encoding a  
15 selected chimeric protein can readily be determined based upon the genetic code for the amino acid sequence of the selected chimeric protein, and, clearly, many nucleic acids will encode any selected chimeric protein. Modifications to the nucleic acids of the invention are also contemplated, since mutations in the steroid receptor binding can thereby be studied for binding affinity. Additionally, modifications that can be useful are  
20 modifications to the sequences controlling expression of the chimeric protein to make production of the chimeric protein inducible or repressible upon addition to the cells of the appropriate inducer or repressor. Such means are standard in the art (*see, e.g.*, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989). The nucleic acids can be  
25 generated by means standard in the art, such as by recombinant nucleic acid techniques, as exemplified in the examples herein, and by synthetic nucleic acid synthesis or in vitro enzymatic synthesis.

          An example of a nucleic acid of the present invention is a nucleic acid encoding  
30 a chimeric protein comprising a green fluorescent protein fused via ten amino acid gly-ala linker to the N-terminus of the rat glucocorticoid receptor. One nucleic acid

encoding this nucleic acid is set forth in SEQ ID NO: 1. This nucleic acid encodes a modified *Aequorea victoria* green fluorescent protein and a modified rat glucocorticoid receptor. Another example is a nucleic acid encoding a chimeric protein comprising a modified *Aequorea victoria* green fluorescent protein fused (via ten amino acid gly-ala linker) to the N-terminus of the human estrogen receptor.

Additionally contemplated by the invention are closely related receptors and nucleic acids encoding them. Thus, provided by the invention are nucleic acids that specifically hybridize to the nucleic acids encoding the chimeric proteins under sufficient stringency conditions to selectively hybridize to the target nucleic acid. Thus, nucleic acids for use, for example, as primers and probes to detect or amplify the target nucleic acids are contemplated herein. Typically, the stringency of hybridization to achieve selective hybridization is about 5°C to 20°C below the T<sub>m</sub> (the melting temperature at which half of the molecules dissociate from its partner). Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA hybridizations. The washing temperatures can similarly be used to achieve selective stringency, as is known in the art. (Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989; Kunkel et al. *Methods Enzymol.* 1987:154:367, 1987).

20

The present invention provides cells containing a nucleic acid of the invention. A cell containing a nucleic acid encoding a chimeric protein typically can replicate the DNA and, further, typically can express the encoded protein. The cell can be a prokaryotic cell, particularly for the purpose of producing quantities of the nucleic acid, or a eukaryotic cell, particularly a mammalian cell. The cell is preferably a mammalian cell for the purpose of expressing the encoded protein so that the resultant produced protein has mammalian protein processing modifications. Additionally, as described above, the cell can have an array of a nucleic acid to which the encoded chimeric protein binds.

30

Labeled compounds and nucleic acids encoding chimeric proteins can be delivered into cells by any selected means, in particular depending upon the purpose of the delivery of the compound and the target cells. Many delivery means are well-known in the art. For example, electroporation, calcium phosphate precipitation, microinjection, cationic or anionic liposomes, and liposomes in combination with a nuclear localization signal peptide for delivery to the nucleus can be utilized, as is known in the art. In particular for transfer of a nucleic acid into a cell, to enhance transfer a cotransfection of the nucleic acid with a second nucleic acid encoding a selectable marker can be performed, and transfected cells selected for by the selectable marker. For example, the interleukin 2 receptor (IL2R) gene can be cotransfected, and selection performed by using beads having the antibody to IL2R bound to the beads to separate out transfected cells. Such methods are standard in the art.

Nucleic acids of the present invention can be used to generate transgenic animals in which the nucleic acid encoding a selected chimeric protein, such as GFP-GR for GR studies or GFP-ER for ER studies, is added to the germ line of the animals. Thus a cell of the invention containing an nucleic acid of this invention is contemplated to include a cell in a transgenic animal. With such transgenic animals, cytoplasm-to-nucleus translocation and gene targeting can be observed in any tissue of interest. Thus studies over the life cycle of the animal can be conducted, so that, for example, development and effects of environment, aging, cancer, etc. can be readily observed. Transgenic animals are generated by standard means known to those skilled in the art.

The provision of the present method to visualize physiologically relevant target sites within the eukaryotic nucleus allows one to directly observe nuclear target sites for any desired steroid- or ligand-dependent transcription factors as well as any nuclearly targeted trans-regulatory factors. Conceptually, the requirements are: 1) to tag the protein of interest with a fluorescent protein, such as the green fluorescent protein, using standard recombinant DNA techniques wherein the two cDNAs are fused in frame to each other such that introduction back into the mammalian cell would give rise to the synthesis of the chimeric protein of interest; and, 2) to create an identifiable nuclear

target site by linking together multiple copies of the potential target site to generate a large enough array as to be readily discernible when the fluorescently labeled chimeric protein interacts with the target site. Such an interaction is visible as an intense, concentrated fluorescent signal unique to the cell harboring the array and absent from the parental cell lacking the array. Agents which modify the interaction of the labeled factor with its cognate site can thus be readily screened.

The present invention further provides a method of screening for a compound that binds to a selected nucleic acid comprising:

- 10       a.     contacting compound fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of the nucleic acid in an array such that the nucleic acid can be directly detected when bound by fluorescently labeled compound; and
- b.     directly detecting the location of fluorescence within the cell,
- 15     fluorescence aggregated at the site of the nucleic acid array indicating a compound that binds to the selected nucleic acid. Thus, the present method can be utilized to directly determine whether a compound binds a nucleic acid, directly or indirectly. For example, transcription factors that have been indirectly shown to affect binding can now be assessed to determine if they do bind the target DNA. If, by this method a factor labeled
- 20     by a fluorescent protein, as herein described (for example, *Aequorea* GFP), is added to cells, and upon visualization of the fluorescence, the fluorescence is seen to be aggregated at the site of the target array of nucleic acid, then the factor, or a factor(s) to which the labeled factor binds, binds the nucleic acid. The nucleic acids are conceptualized as merely serving as an easily identifiable "stage" upon which the
- 25     "actors" (i.e., transcription factors and associated factors) appear.

In the present inventive methods, by fluorescence aggregated at the site of the nucleic acid array is meant fluorescence aggregated at a single predominant site within the nucleus. Such aggregation can readily be detected upon excitation of the fluorescent protein. Detection, as described above, can be performed with the use of a fluorescent

30     microscope. Magnification can be altered as desired for more or less detail in visualizing

the aggregation, such as from about 60x to about 200x, with a typical magnification being about 100x. The fluorescent protein in any herein described method can be, for example, a green fluorescent protein, as described herein.

5           Compounds for use in the present methods can be labeled by standard means in the art for linking a peptide to the compound. For example, when the compound to be labeled is a peptide, polypeptide or protein, a chimeric protein can be made by synthesizing a nucleic acid that encodes the chimeric protein having the fluorescent protein fused to the compound protein. Proteins can also be labeled with a fluorescent  
10 protein by a chemical bridge. Additionally, a fluorescent protein label can be placed on the compounds by non-covalent interaction, such as that displayed by steroids with their binding domain [e.g., fluorescein or rhodamine conjugated dexamethasone (available through Molecular Probes, Inc. catalog number D-1382 or D-1383) for the glucocorticoid receptor].

15           More than one chimeric protein, preferably each comprising a fluorescent protein that emits a different wavelength of light (for example a modification of *Aequorea* GFP that emits blue and a modification of *Aequorea* GFP that emits green), can be used simultaneously in the present inventive methods. For example, plasmids pCI-nGL2-C656G;  
20 C656G; pCI-nGL3-C656G; pCI-nGL4-C656G; pCI-nGL5-C656G; pCI-nGL7-C656G; pCI-nGL9-C656G; pCI-nGL10-C656G; pCI-nGL11-C656G, all GFP-GR having blue variants of GFP (BFP) in a humanized codon usage (improved blue variants) can be used with any GFP-GR having green GFP (such as pCI-nGFP-C656G, pCI-nGL1-C656G, pTET-nGFP-C656G and pOP-nGFP-C656G, exemplified herein) in  
25 this method.

          Alternatively, two different fluorescent moieties that show distinct excitation maxima with identical emission wavelengths could also be simultaneously utilized to detect the presence of both fluorescently labeled compounds on the same DNA array.  
30 With the appropriate combination of fluorescent molecules only one excitation wavelength may be required to detect the presence of both fluorescent moieties in close

proximity through the process of fluorescent energy transfer, wherein the excitation wavelength excites one moiety which emits at the absorption wavelength of the second moiety. Such simultaneous use will allow the detection of interaction of various transcription factors and cofactors with each other and with the DNA to activate and/or  
5 repress transcription from a specific regulatory sequence.

Cells for use in the present methods are cells having a plurality of copies of the nucleic acid in an array such that the nucleic acid can be directly detected when bound by fluorescently labeled compound. Such cells are described herein and can be prepared  
10 as described herein. An example of such a cell is cell line 3134, having about two hundred copies of the MMTV LTR-Ha-v-ras- gene, each of which has four copies of the binding site for glucocorticoid receptor, in direct tandem repeats integrated into the genome of the cell. Cells further can have a nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to the binding compound/transcription factor of  
15 interest, *i.e.*, the binding compound/transcription factor which may bind the nucleic acid in the array in the cell. For example, 3134 cells containing pCI-nGFP-C656G, pCI-nGFP-C656G, pTET-nGFP-C656G or pOP-nGFP-C656G are exemplified herein. Nucleic acids encoding a chimeric protein can either be integrated or not, as best suits the specific method being performed.

20

The present invention also provides a method of characterizing cells in which a compound fluorescently labeled by a fluorescent protein is expressed in the cell and the localization of the fluorescent protein observed for perturbation in localization of the fluorescently labeled protein in the absence or presence of signals that affect protein  
25 function (example of GFP-ER in the MCF7 and MDA-MB-231 cell line).

The present invention also provides a method of screening for a ligand that activates gene targeting of a steroid receptor in the nucleus of a mammalian cell comprising contacting the ligand with a cell having a plurality of steroid receptor  
30 response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a

nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and directly detecting the location of fluorescence within the cell, fluorescence aggregated at the site of the steroid receptor response element array in the nucleus indicating a ligand that activates the gene targeting of a steroid receptor in the nucleus of a mammalian cell. A ligand for any steroid receptor can be determined by this method by creating an array of the steroid receptor response element in the cell used. For example, cell line 3134 can be used to detect ligands that activate gene targeting of glucocorticoid response element, progesterone receptor, or aldosterone receptor. An example of a chimeric protein for use in this method in, for example cell line 3134, is one that has a green fluorescent protein, such as *Aequorea* fluorescent protein, fused to the N-terminus of glucocorticoid receptor, such as the chimeric protein comprising the amino acid sequence set forth in SEQ ID NO: 2.

The present invention further provides a method of screening for a ligand that activates the translocation of a steroid receptor to the nucleus or redistribution of a steroid receptor in a mammalian cell comprising contacting the ligand with a cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and directly detecting the location of fluorescence within the cell, change in the location of the fluorescence (*e.g.*, cytoplasm to nucleus in the case of the glucocorticoid receptor or redistribution within each cellular compartment; *e.g.*, for primarily nuclear receptors, such as the progesterone receptor, aggregation on the MMTV LTR-array in the 3134 cell) could indicate a potential ligand of the receptor. In the present method, redistribution of the receptor can be directly observed. Also, in the case of the glucocorticoid receptor, translocation to the nucleus, even in the absence of binding to the nuclear DNA, can be seen; and importantly, one can see if a ligand causes only translocation to the nucleus (by location of fluorescence primarily in the nucleus, but in a diffuse or reticular, rather than aggregated, pattern) or causes both translocation to the nucleus and binding to nuclear DNA (by location in the nucleus aggregated primarily at a site). In the case of estrogen

receptor which has been partially activated due to trace estrogenic substances present in the culturing media, the receptor is nuclearly localized but shows dramatically different intranuclear distribution in two human breast cancer cell lines. In the case of MCF7 cell, a human adenocarcinoma breast cell line which contains endogenous estrogen receptor and shows hormone dependent growth, the GFP-tagged estrogen receptor is collected upon nuclear structures and shows focal accumulation patterns. In contrast, the MDA-MB-231 cell, a human adenocarcinoma breast cell line which lacks endogenous estrogen receptor and shows hormone-independent growth, the GFP-tagged estrogen receptor

under the same culturing condition is extremely diffuse. These two dramatic differences in the nuclear localization patterns in two different human breast cancer cell lines suggests potential usefulness of GFP-ER as a diagnostic reagent for characterizing different human breast cancer cells as well as characterizing the progression of human breast cancer. These

differences in GFP-ER localization patterns also suggest an additional requirement for cellular components in permitting the targeting of the estrogen receptor onto nuclear structures; these cellular components maybe absent as the cell progresses from hormone-dependent to a hormone-independent stage in the progression of human breast cancer.

20

The present invention additionally provides a method of detecting in a biological sample the presence of an agonist of a steroid receptor comprising contacting the sample with a cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and directly detecting the location of fluorescence within the cell, the location of fluorescence aggregated at the site of the steroid receptor response element array in the nucleus indicating the presence of an agonist of the steroid receptor in the sample.

30



Biological samples can include any relevant sample from the body, such as blood, plasma, urine and saliva.

The present invention also provides a method of detecting in a biological sample  
5 the presence of an antagonist of a steroid receptor comprising contacting the sample and both agonist and antagonist of the steroid receptor with a cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent  
10 protein is fused to the steroid receptor; and directly detecting the location of fluorescence within the cell, the absence of fluorescence substantially aggregated at the site of the steroid receptor response element array in the nucleus indicating the presence of an antagonist of the steroid receptor in the sample.

Also provided is a method of monitoring the level of an agonist of a steroid  
15 receptor in a subject comprising periodically obtaining a biological sample from the subject, contacting the sample with a cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid  
20 encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor; and directly detecting the location of fluorescence within the cell, a decrease in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating a decrease in level of the steroid agonist of the steroid receptor in the sample and an  
25 increase in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid agonist of the steroid receptor in the sample.

Further, provided by the present invention is a method of monitoring the balance  
30 between levels of an agonist of a steroid receptor and an antagonist of the steroid receptor in a subject comprising periodically obtaining a biological sample from the

subject, contacting the sample with a cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid  
5 receptor; and directly detecting the location of fluorescence within the cell, an increase in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid agonist relative to level of the steroid antagonist in the sample, and a decrease in fluorescence aggregated at the site of the steroid receptor  
10 response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid antagonist of the steroid receptor relative to level of the steroid agonist in the sample.

The present invention also provides a method of determining an effective dosage  
15 of a steroid receptor agonist in a subject comprising transferring into a set of cells from the patient a nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a steroid receptor; contacting the cells in the set with one of a selected range of dosages of the steroid agonist; and directly detecting location of fluorescence in the set of cells, a dosage capable of locating fluorescence substantially in the nucleus indicating  
20 an effective dosage of steroid receptor agonist.

Further provided by the present invention is a method of determining an effective dosage of a steroid receptor agonist to maintain steroid receptor activation for a selected period of time in a subject comprising administering to the subject a dosage of the  
25 steroid receptor agonist; periodically obtaining a biological sample from the subject; contacting the sample with a mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor and the cell further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the  
30 steroid receptor; and directly detecting the location of fluorescence within the cell, a dosage that maintains the location of fluorescence at the site of the steroid receptor

response element array in the nucleus for the selected period of time indicating an effective dosage.

The present invention additionally provides a method of detecting a defect in a response pathway of a steroid receptor in a subject comprising transferring into a cell  
5 from the subject a nucleic acid functionally encoding a chimeric protein comprising a fluorescent protein fused to the steroid receptor and detecting the location of fluorescence within the cell as compared to the location of fluorescence within a normal, control cell transfected with the nucleic acid, a difference in location of fluorescence  
10 within the cell of the subject as compared to location of fluorescence within the normal, control cell indicating a defect in the response pathway of the steroid receptor.

The present invention also provides a method of determining whether a defect in a response pathway of a steroid receptor in a subject is in translocation of the steroid  
15 receptor to a cell nucleus, comprising transferring into a cell from the subject having the defect a nucleic acid functionally encoding a chimeric protein comprising a fluorescent protein fused to the steroid receptor and detecting the location of fluorescence within the cell, the location of fluorescence substantially in the cytoplasm of the cell indicating the defect is in translocation of the steroid receptor to the nucleus.

20

The present invention also provides a method of characterizing a ligand's effect on cellular localization of a compound to which the ligand binds in a cell comprising contacting the ligand with a cell having the compound fluorescently labeled by a fluorescent protein and directly detecting the location of fluorescence within the cell, the  
25 location of fluorescence in the cell indicating the localization effect of the ligand on the compound. Compounds can be, *e.g.*, steroid receptors, transcription factors and the like. For example, the examples provide characterization of localization of GR in response to two ligands, dexamethasone and RU486, and characterization of the localization of ER in response to ligands, agonist beta-estradiol or anti-estrogens,  
30 4-hydroxytamoxifen or ICI164384." By this method, the ligands triggering binding of so-called "orphan receptors" to their binding site(s) can be discovered.

Additionally provided is a method of determining a binding site for a DNA-binding protein comprising contacting the DNA-binding protein fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of a nucleic acid having a putative binding site in an array such that the putative binding site can be directly  
5 visualized when bound by the fluorescently labeled DNA-binding protein, and directly detecting the location of fluorescence within the cell, the presence of fluorescence aggregated at the putative binding site indicating a binding site to which the DNA-binding protein binds. The absence of fluorescence aggregated at the putative binding site can suggest a binding site to which the DNA-binding protein does not significantly  
10 bind.  
the location of fluorescence in the cell indicating the localization effect of the ligand on the compound.

Also provided by the present invention is a method for screening for gene-  
15 specific combinations of compounds that bind the gene specifically, comprising contacting (a) a first compound labeled by a fluorescent protein emitting a first spectrum of light and a second compound labeled by a fluorescent protein emitting a second spectrum of light with (b) a cell having a plurality of copies of the regulatory region of the gene in an array such that the regulatory region can be directly detected when bound  
20 by compound labeled by a fluorescent protein. Fluorescence for the first and second spectrum is then localized. Aggregation of fluorescence of both the first and second spectrum at the site of the regulatory region array would indicate a combination of compounds that binds the gene specifically; the location of only one spectrum aggregated at the array would indicate that only the corresponding compound binds the  
25 gene of interest directly. By using a combination of screens, compounds that bind the specific DNA both directly and indirectly can be determined for a gene of interest. Such gene-specific combinations of compounds can be used to develop gene-specific drugs that interfere with transcription activators in a selective manner. This method is based on the fact that each transcription factor, cofactor, etc. affects many genes, but for each  
30 gene there is likely only one combination of factors and cofactors that activates/represses it. Therefore, once, by this screening method, it has been determined

which combination of factors causes activation or repression of a specific gene, then a combination of drugs, to affect all relevant factors for that specific gene, can be administered to selectively activate/repress that gene. Thus a combination of drugs can ultimately be used to activate or repress the selected gene.

5

#### Statement Concerning Utility

The present invention provides methods for directly detecting the binding of compounds to nucleic acids. The present invention allows a simple and straightforward manner in which direct interaction between a sequence-specific DNA binding protein or  
10 its co-factor and its putative regulatory site in the *in vivo* genomic context can be addressed. For example, GFP-steroid fusion proteins and cell lines containing receptor binding sites (response elements) in multimerized arrays are provided for direct visualization of *in vivo* gene targeting. These reagents provide a simple, rapid, straightforward, sensitive, and biologically relevant assay for each target nucleic acid  
15 and binding compound. These reagents can in turn be used for several medically important applications, including diagnostic tests for concentration of cognate ligand in clinical samples (urine, saliva, blood, etc.). Additionally, the reagents can be used for direct tests for defects in steroid pathways in subjects. The reagents further can be used for screening of chemical banks for compounds with ligand agonist activity for each  
20 receptor, and development of drugs based on these activities. Furthermore, using the fluorescent protein-steroid receptor fusions, colocalization of a selected receptor with any other cofactor that may be recruited to the chromosome template can be evaluated. Using a separate tag (different color) for the DNA target, the steroid receptor can be fused to any factor that may be recruited by the steroid receptor and determine if the  
25 factor colocalizes on the DNA target when the steroid receptor is activated, allowing for a direct test for defects in factor colocalization in human disease/syndromes. This ability to observe direct interaction of any trans-regulatory factor or co-factor and its regulatory site *in vivo* provides a screening method for useful and novel drugs, directed against trans-regulators, and the development of gene-specific multi-drug therapies.  
30 Using the glucocorticoid receptor as an example and known ligands of the receptor (e.g., agonist dexamethasone and antagonist RU486), the validity of the methodology is

herein demonstrated. In a specific example, the ability to monitor effective hormone concentration by this novel methodology in real time can lead to the development of diagnostic kits that can be used to properly gauge the required amounts of hormone administered to patients requiring long-term or short-term hormone treatment. In addition, in the case of those trans-acting factors regulated at the level of the nuclear/cytoplasmic and cytoplasmic/nuclear translocation step, diseases arising from such a failure can be directly diagnosed by fusion of the regulatory molecule to a fluorescent moiety using standard recombinant DNA techniques. With this simple, inventive methodology, novel classes of drugs directed not only against members of the steroid-ligand-dependent transcription factors but to new classes of drugs that target other transcription factors or their co-factors can be screened. Furthermore, by using combination of drugs which target certain trans-regulatory factors either specifically or selectively, a gene-specific based drug therapy regimen can be created. This multi-drug therapy designed against a certain critical gene implicated for a particular human disease would be tailored to affect the activity of the trans-regulatory factors all of which act synergistically to regulate the transcription of the gene implicated in the disease. Furthermore, the reagents allow the development of transgenic animals containing fusion proteins such as each of the GFP-receptor fusions that can provide a unique tool to study subcellular distribution of the receptors in all tissues of the animal, and the effect of pharmacologic agents on function of each of the receptors in the various tissues. Numerous other utilities will be apparent to the skilled artisan in light of the present invention.

## EXAMPLES

The present invention is more particularly described in the following examples which are intended as illustrative only since numerous modifications and variations therein will be apparent to those skilled in the art.

### 30 Plasmids.

- pCI-nGFP-C656G:** Plasmid pCI-nGFP-C656G was derived from pCI-nH6HA-C656G (Smith et al., submitted) and pZA69 (a kind gift from Mike Moser and Ravi Dahr). pZA69 contains a BspHI fragment of pZA66, a plasmid containing S65T GFP (Tsien, R.Y. (1995) *Nature* 373,663-664) with the internal NcoI site removed by a silent mutation. The pCI-nH6HA-C656G DNA expresses the rat glucocorticoid receptor with the C656G mutation (kindly provided by S. Simons, Jr.) under the control of the CMV promoter/enhancer, and is tagged at the N-terminus with (his)<sub>6</sub> and hemagglutinin epitope recognized by monoclonal antibody 12CA5 (Niman, H. L., Houghten, R. A., Walker, L. E., Reisfeld, R. A., Wilson, I. A., Hogle, J. M. & Lerner, R. A. (1983) *Proc. Natl. Acad. Sci. U. S. A.* 80, 4949-4953); this DNA was cleaved at a unique site with PvuII, separating the two tags from the rest of the glucocorticoid receptor. A 768 bp DNA fragment containing the GFP cDNA with the S65T mutation was inserted at the PvuII site. This GFP DNA fragment is obtained by digesting pZA69 with BglI, attaching a BglI hairpin linker with the sequence (5'-GCGCGCTGATCAGAATTCCTTTTAGGAATTCTGATCAGCGCGCTGA-3') (SEQ ID NO:3), recutting the resulting DNA with BclI and XhoI, and then filling-in with the large fragment of DNA polymerase (Klenow) to create a 768 bp blunt-end fragment.
- pCI-nGL1-C656G:** The GFP (S65T) variant with the jellyfish codon usage in the mammalian expression GFP-GR vector, pCI-nGFP-C656G, has been replaced with a humanized codon usage from the plasmid, pGreenLantern-1 (LifeTechnologies, Inc., Gaithersburg, MD, catalog number 10642-015), for improved translation in mammalian cells [this humanized GFP is also approved for in vitro diagnostic use by LifeTechnologies] to generate an improved mammalian expression GFP-GR plasmid, pCI-nGL1-C656G.
- pCI-nGL2-C656G; pCI-nGL3-C656G; pCI-nGL4-C656G; pCI-nGL5-C656G; pCI-nGL7-C656G; pCI-nGL9-C656G; pCI-nGL10-C656G; pCI-nGL11-C656G: Blue variants of GFP (BFP) in a humanized codon usage (improved blue variants) have been generated which are fused to the rat glucocorticoid receptor (C656G). Each has a

different chromophore. Site-directed mutagenesis of the GFP element of pCI-nGL1-C656G (which has humanized GFP) was performed to introduce known chromophore-altering mutations into the GFP element. Site-directed mutagenesis was performed using Chameleon™ double-stranded site-directed mutagenesis kit (Stratagene, catalog number 200509). These new fusion plasmids were then expressed in 1471.1 cells. Fluorescent spectrophotometry indicates the chromophores are present and that the GFP-GR fusion is intact. Additional mutants, such as those producing a longer wavelength chromophore, *e.g.* red chromophores, can be made by the same methods.

10       **GFP-ER plasmid:** The glucocorticoid receptor portion of pCI-nGL1-C656G DNA has been replaced with a human estrogen receptor (ER). ER binds to an ER response element in cells. The GFP-ER was found to be functional in both transcriptional activation as well as proper subcellular localization in several cell lines, as described below. GFP-ER plasmid is made using the same site-directed mutagenesis as  
15 for preparing blue variant plasmids. Briefly, a cDNA encoding ER is inserted into pCI-nGL1-C656G in place of the GR cDNA. An ER cDNA (Green, Stephen, *et al.*, *Nature* 320:134-139 (1986) (GenBank accession number X03635) (note: this sequence has a Gly<sub>400</sub> to Val<sub>400</sub> mutation); Greene, Geoffrey L., *et al.* *Science* 231:1150-1154 (1986) (GenBank accession number M12674 (having Gly<sub>400</sub> to Val<sub>400</sub> mutation)); Pfeffer, U.  
20 *Cancer Res.* 53:741-743 (1993) (GenBank accession number X73067) (ER fragment having correct Gly<sub>400</sub> region coding sequence)) is mutagenized to create a MluI site at the start point of translation of ER and a SalI site in the 3' untranslated region of the cDNA (alternatively, the ER sequence can be generated by PCR). pCI-nGL1-C656G has a unique BssHII site after the (gly-ala)<sub>5</sub> linker and a SalI site after the GR portion of  
25 the plasmid DNA. The DNA cut end made by MluI is complementary to the end made by the BssHII. Therefore, the GR cassette can be removed by BssHII/SalI digestion and the ER cassette (released by MluI/SalI digestion) subcloned into the remaining vector at the BssHI and SalI cut ends.

30       **pOP-nGFP-C656G:** The original cDNA encoding GFP-GR from the plasmid, pCI-nGFP-C656G, has been subcloned into a tetracycline-regulatable mammalian



expression vector, pTET-Splice (Life Technologies, catalog number 10583-011), to give pTET-nGFP-C656G, and a lac-regulatable mammalian expression vector, pOPRSVI CAT (Stratagene Cloning Systems, catalog number 217450) to give pOP-nGFP-C656G.

- 5           Other plasmids used in this study are: pLTRLuc (full-length MMTV LTR driving the expression of a luciferase gene) (Lefebvre, P., Berard, D. S., Cordingley, M. G. & Hager, G. L. (1991) *Mol. Cell. Biol.* **11**(5), 2529-2537), pCMVIL2R (IL2R expression plasmid) (Giordano, T., Howard, T. H., Coleman, J., Sakamoto, K. & Howard, B. H. (1991) *Exp Cell Res.* **192**, 193-197), and pUC18 (Life Technologies,  
10   Inc.).

- GFP-fusion plasmids of the invention are tested for expression and subcellular localization by transfection into several mammalian cells. For example, GFP-GR plasmids were analyzed in C127 (mouse) cells, HeLa (human cervical cancer) cells, and  
15   MCF7 (human breast adenocarcinoma) cells (ATCC accession number HTB22). GFP-ER plasmids were analyzed in 1471.1 cells, C127 (mouse) cells, MCF7 (human breast adenocarcinoma) cells, and MDA-MB-231 (human breast adenocarcinoma) cells (ATCC accession number HTB 26). Localization is observed in the absence of added  
20   hormone in either 5% or 10% charcoal-stripped fetal calf serum. Cells having GFP-fusion plasmids are then treated with a selected ligand, and subcellular localization and quantitative observations are made.

#### **Cell Line 1471.1 and derivatives**

- Cell line 1471.1 contains multiple copies of a BPV MMTV-LTR-  
25   chloramphenicol acetyltransferase (CAT) reporter gene fusion introduced in the murine adenocarcinoma C127 cell (Archer, T. K., Cordingley, M. G., Marsaud, V., Richard-Foy, H. & Hager, G. L. (1989) in *Proceedings: Second International CBT Symposium on the Steroid/Thyroid Receptor Family and Gene Regulation*, eds. Gustafsson, J. A., Eriksson, H. & Carlstedt-Duke, J. (Birkhauser Verlag AG, Berlin),  
30   pp. 221-238).

Derivatives of 1471.1 cells which contain over a thousand copies of the MMTV LTR-CAT have also been generated with the tetracycline- and lac-regulatable GFP-GR (pTET-nGFP-C656G and pOP-nGFP-C656G), e.g., cell line 3677. In these derivative cell lines, GFP-GR expression occurs upon tetracycline withdrawal from 5 ug/ml or  
5 upon induction with IPTG using standard procedures and as recommended by the manufacturer of these inducible systems.

**Transfection.** Plasmid DNA was transiently introduced into 1471.1 cells either by calcium phosphate coprecipitation using a BES-based buffer (Chen, C. & Okayama, H.  
10 (1987) *Mol. Cell Biol.* 7, 2745-2752) or by electroporation. For calcium phosphate coprecipitation, semi-confluent cells maintained in Dulbecco's modified eagle media (DMEM; Life Technologies, Inc.) supplemented with 10% fetal calf serum (FCS; Life Technologies, Inc.), 2 mM glutamine, and 50 mg/ml gentamicin sulfate were trypsinized,  
15 washed, resuspended at  $7 \times 10^4$  cells/ml in DMEM supplemented as above except FCS was treated with charcoal/dextran-treated fetal bovine serum (Hyclone Laboratories, Inc.), dispensed as 1 ml into 2x2 cm<sup>2</sup> Lab-Tek Chamber Slide (Nunc, Inc.) or as 10 ml into a 100 mm petri dish layered with 24.5 mm-diameter Dvorak-Stotler coverslips (Nicholson Precision Instrument). Cells were grown overnight in a 37°C humidified incubator with 5% CO<sub>2</sub>. The following morning, media was replaced with fresh  
20 supplemented DMEM containing dextran/charcoal treated FCS, and in the afternoon, cells were transfected with 1 ml of transfection mixture containing 20 µg plasmid DNA (as indicated in the appropriate figure legend) per 10 ml of cells, essentially as described (Chen, C. & Okayama, H. (1987) *Mol. Cell Biol.* 7, 2745-2752). Cells were left overnight in a 37°C humidified incubator with 2.9% CO<sub>2</sub>. About 12-16 hours after  
25 transfection, media was replaced, and the cells were allowed to recover for two hours before further treatment and imaging. For calcium depletion experiments, cells were electroporated with 5-20 µg pCI-nGFP-C656G DNA for  $2 \times 10^7$  cells in 0.2 ml cold DMEM at 250 V and 800 µF, left to recover on ice for 5 minutes, and then diluted in DMEM supplemented with dextran/charcoal treated FCS before plating. Cells were  
30 then grown for 12 to 16 hours in a 37°C humidified incubator at 5% CO<sub>2</sub>. Before treatment and imaging, cells were fed with fresh media.

**Enrichment of Transfected Cells and Analysis of Cytosolic Extracts.** Cells that took up exogenous DNA were enriched by cotransfection with pCMVIL2R, an IL2R (interleukin 2 receptor) expression plasmid, and selection for IL2R<sup>+</sup> cells using magnetic beads (Dynal) coated with mouse anti-human IL2R antibody (Boehringer Mannheim, clone 3G10), as described (Giordano, T., Howard, T. H., Coleman, J., Sakamoto, K. & Howard, B. H. (1991) *Exp Cell Res.* **192**, 193-197). Extracts from the IL2R<sup>+</sup> and IL2R<sup>-</sup> cells were made by three cycles of freezing and thawing of the cell suspension in either 100 mM sodium phosphate (pH 7.8) with 1 mM DTT or 250 mM Tris-HCl (pH 7.8). After clarifying the lysate, extracts made with the phosphate buffer was used to assay for the amount of luciferase activity in a MicroLumat LB96P as recommended by the manufacture, EG&G Berthold. For the Tris-buffered extract, CAT activity was assayed as described (Gorman, C. M., Moffat, L. F. & Howard, B. H. (1982) *Mol. Cell Biol.* **2**, 1044-1051). Protein concentration was determined by the method of Bradford using the Bio-rad Protein Assay reagent (Bio-rad Laboratories, Inc.).

**Determination of Intracellular Calcium.** Intracellular free calcium concentrations were determined in single cells by measuring the signal from the calcium sensitive indicator Fura-2, according to Tsien and Harootunian (Tsien, R. Y. & Harootunian, A. T. (1990) *Cell Calcium* **11**, 93-109). Briefly, cells were cultured on cover slips and electroporated with the GFP-GR chimera one day before microscopy. In preparation for imaging, cells were treated for 30 min with either assay buffer (Hank's balanced salt solution without phenol red, with 2 mg/ml glucose and 1 mg/ml BSA, containing 3 mM Ca<sup>++</sup>) or with calcium-free buffer (Eagle's No. 2 medium without calcium, containing 1 mg/ml BSA, 5 mM EGTA, 5  $\mu$ M thapsigargin, 2  $\mu$ M ionomycin). The cells were then loaded with 5  $\mu$ mol/L Fura-2-AM (from Molecular Probes Inc.) and 0.02% pluronic F-127 with either calcium-free or calcium supplemented media (30 min at room temperature, washed three times, then incubated for 15 min at 37°C). After loading, cells were placed into a Dvorak-Stotler chamber (inner volume 224  $\mu$ l) and perfused at 37°C with either calcium-containing or calcium-free media. Intracellular

calcium content was measured in three independent experiments—at least 20 cells in each experiment. Ratio imaging was performed using Image 1 software (Universal Imaging Corp.) running on an IBM PC, using 340 nm and 380 nm excitation, 510 nm emission, and 490 nm dichroic barrier filters, a Zeiss Photomicroscope III microscope, enclosed into a temperature controlled incubator, and an intensified (Videoscope) CCD camera (Dage 72), and optical disc recorder (Panasonic). The system was calibrated for  $[Ca^{2+}]_i$  measurement using Fura-2 pentapotassium salt and calibration buffer kit from Molecular Probes Inc. Intracellular free calcium concentrations in cells with calcium supplemented buffer were  $350 \pm 183$  nM, while in calcium-free buffer  $60 \pm 11$  nM.

10

**Image Acquisition and Analysis.** For time course studies, cells were placed into a Dvorak-Stotler chamber (inner volume 224  $\mu$ l) and perfused at 37°C with assay buffer for 3 min, then with the same buffer containing 1 nM dexamethasone (dex) for 2 hours at 10 ml/hour flow rates. Samples were evaluated using a Zeiss Axiovert 10 microscope surrounded by an incubator and equipped for epifluorescence with illumination from XBO burner, 480 nm excitation and 535 nm emission and 505 nm dichromatic barrier filters (from Chroma Technology Corp.). Images were acquired every 15 seconds with a high resolution, cooled CCD camera equipped with an electromechanical shutter (Photometrics p200). Images were collected on Silicon Graphics workstations (4D310-VGX), using custom software, incorporating functions from a vendor supplied library (G.W. Hannaway & Assoc.). Experiments requiring real time image acquisition were performed on the imaging system described for the intracellular calcium measurements.

25

Confocal laser scanning microscopy was carried out on a Nikon Optiphot microscope equipped with BioRad MRC-600 confocal laser scanning unit, with fluorescent excitation produced by the 488 nm line of a krypton-argon laser, and using a Fluor 100/1.3 oil phase objective. From living cells expressing GFP-GR, serial 0.5  $\mu$ m optical sections were collected and the digitized images were imported into a Silicon Graphics Indigo 2 workstation. Three dimensional image rendering, analysis and reconstruction was carried out with the ANALYZE software from the Mayo Clinic.

30

**Tagging of a Highly Dexamethasone-Sensitive Form of GR with a Highly Fluorescent Variant of GFP.** To develop a highly efficient, fluorescent version of the glucocorticoid receptor, we generated a GFP-GR chimera in which cDNA encoding a 27 kDa GFP variant is fused in frame to the second amino acid of a rat glucocorticoid receptor (Fig. 3A). The GFP variant contains a serine to threonine substitution at amino acid 65 (S65T mutation) from the jellyfish *Aequorea victoria*, which increases the efficiency of formation of the GFP chromophore by accelerating the rate of oxidation required for chromophore generation. In addition, the resulting chromophore is six-fold more fluorescent than the wild-type GFP (Heim, R., Cubitt, A. B. & Tsien, R. Y. (1995) *Nature* **373**, 663-664), making the use of this chromophore perhaps the most sensitive method for labelling proteins (Wang, S. & Hazelrigg, T. (1994) *Nature* **369**, 400-403). Additionally, the chromophore is formed faster, potentially explaining why expression at 37°C in mammalian cells is achieved herein, contrary to reports that the GFP chromophore does not form at a relatively high temperature of 37°C (Ogawa, *et al.* *Proc. Natl. Acad. Sci. USA* **92**:11899-11903 (1995)).

Since glucocorticoid receptor is ubiquitously present in all mouse cells and selective activation of the tagged receptor is required to assess the functionality of the receptor, we therefore used a glucocorticoid receptor having a higher affinity for its ligand than the endogenous receptor. To this end, S65T GFP was fused to a rat glucocorticoid receptor that contains a cysteine to glycine mutation at position 656 of the steroid binding domain (Chakraborti, P. K., Garabedian, M. J., Yamamoto, K. R. & Simons, S. S. J. (1991) *J. Biol. Chem.* **266**, 22075-22078). This point mutation, C656G, increases the affinity of the receptor ten-fold for its ligand. A dose response curve shows complete activation of GFP-GR at 1 nM dexamethasone and half maximum at 0.1 nM; the endogenous mouse receptor is fully activated at 100 nM dexamethasone with the half maximal stimulation at 10 nM. Thus, presence of the C656G mutation permits selective activation of the transfected chimeric receptor without activation of the endogenous receptor.

**Transcriptional Competence of GFP-GR.** When the plasmid encoding this chimera, pCI-nGFP-C656G, is introduced into cultured mouse cells, a fusion polypeptide with the predicted molecular weight of 118 kDa is produced. When these cells are stimulated with 1 nM dexamethasone, a co-transfected reporter construct (pLTRLuc) containing the luciferase reporter gene under the control of the mouse mammary tumor virus promoter (MMTV LTR) is activated (Fig. 3B). In addition, 1 nM dexamethasone-treated cells show accumulation of luciferase activity dependent on the amount of GFP-GR expression plasmid included in the transfection. In the absence of any GFP-GR expression plasmid, no significant luciferase activity accumulated in the 1 nM dexamethasone-treated cells, indicating that 1 nM dexamethasone activated the GFP-GR chimeric protein but not the endogenous GR. With increasing amount of the GFP-GR expression plasmid, luciferase activity in the 1 nM dexamethasone-treated IL2R<sup>+</sup> cells reaches the same level as that in 100 nM dexamethasone-treated IL2R<sup>+</sup> cells lacking GFP-GR expression plasmid (compare 2 µg pCI-nGFP-C656G, 1 nM dex with 0 µg pCI-nGFP-C656G, 100 nM dex). Since the 100 nM dexamethasone treatment gives complete activation of the endogenous GR in the latter case, we conclude that the GFP-GR chimeric receptor is fully functional in dexamethasone-mediated transcriptional activation of the transiently introduced reporter plasmid DNA.

Furthermore, derivative cell lines of 1471.1 which contain over a thousand copies of the MMTV LTR-CAT have also been generated with the tetracycline- and lac-regulatable GFP-GR (pTET-nGFP-C656G and pOP-nGFP-C656G), allowing GFP-GR expression upon tetracycline withdrawal from 5 µg/ml or upon induction with IPTG using standard procedures and as recommended by the manufacturer of these inducible systems. In these cells, acceptable levels of GFP-GR is reached after overnight induction of the regulatable promoters.

**Assay for Ligand Effects:** GFP-fusion plasmids of the invention are assayed for effects of a selected ligand on subcellular localization by transfection into selected cells. C127 (mouse) cells, HeLa (human cervical cancer) cells, and MCF7 (human breast adenocarcinoma) cells (ATCC accession number HTB22) were each transfected with

GFP-GR plasmids and with GFP-ER plasmids. Localization is first observed in the absence of added hormone in either 5% or 10% charcoal-stripped fetal calf serum. Cells having GFP-fusion plasmids are then treated with a selected ligand, and subcellular localization and quantitative observations are made.

5

**GFP-GR Results:** GR is observed to be localized to the cytoplasm in the absence of added ligand. However, when cells having the GFP-GR plasmids were treated with dexamethasone, foci of fluorescent signal (massive variation in the peak and valley of fluorescent signal) were observed in the nucleus (*see, e.g.*, FIG 4(A)). When  
10 these cells were treated with RU486, the fluorescent signal is near uniform, with a matrix-like appearance where fluorescent signal appears (*see, e.g.*, FIG 4(B)).

**GFP-ER Results:** ER is observed to be localized to the nucleus in the absence of added estrogen hormone. When a hormone-dependent, estrogen receptor positive  
15 human breast cancer cell line, MCF7, is transfected with GFP-ER expression plasmid, expression of GFP-ER results in nuclear signal that is structured with peaks and valley of concentration of signals (FIG. 5(A)). Although the GFP-ER is already partially activated due to trace estrogenic substances present in the culturing medium, addition of  
20 agonist beta-estradiol leads to further concentration of nuclear signals on nuclear structures. In the case of a hormone-independent, estrogen receptor negative human breast cancer cell line, MDA-MB-231, the expressed GFP-ER in the absence of added ligand shows a diffuse, fuzzy pattern, with only hints of attachments to structures in the nucleus.

25

These results provide an example of use of the present method as a diagnostic for missing, dysfunctional or nonfunctional components in any selected cell. For example, upon determining that a ligand, such as estrogen for ER binding to its response element, is present in a cancer cell and used for growth or maintenance of the cell, one  
30 can treat the cancer by administering a compound to deprive the cell of that ligand, such as by administering anti-estrogen to a breast cancer showing the same results as MCF7

(hormone-dependent) to reduce growth of the cells. Similarly, such a diagnostic can tell one if such a treatment in another cell would be futile because the cell is not dependent upon that ligand.

5        These results also demonstrate that the present method can be used as a screen to classify cell types for the ability (or lack of ability) to target nuclear structures, to traffic compounds in a particular pattern, etc. This also provides information for selecting treatment regimens for various diseases or disorders, based on activating, inactivating or altering the function of ligands in the cell.

10

Transcriptional activation of the MMTV LTR target genes by GFP-GR shows a ligand specificity characteristic of glucocorticoid receptor. When activated by dexamethasone, GFP-GR is competent to induce not only the transiently introduced MMTV LTR-luciferase reporter DNA, as mentioned above (Fig. 3B), but also the  
15 multi-copy MMTV LTR-CAT reporter genes present in 1471.1 cells (1 nM and 10 nM dex, Fig. 3C). In contrast, treatment with 10 nM RU486, an antagonist with little GR agonist activity, or progesterone, a poor agonist, results in little activation of the MMTV LTR-CAT reporter; 17- $\beta$ -estradiol, a steroid that shows no affinity for GR, fails to activate the LTR. Thus, the ligand specificity of GR in the transcriptional activation  
20 of the MMTV LTR is maintained in GFP-GR expressing cells.

#### **Visualization of GFP-GR Cytoplasm-to-Nuclear Translocation in a Single**

**Metabolically Active Cell.** Because the S65T variant of the GFP chromophore is resistant to photobleaching (Heim, R., Cubitt, A. B. & Tsien, R. Y. (1995) *Nature* 373,  
25 663-664), it was possible to use confocal and time-lapse video microscopy to observe GFP-GR over extended periods. Using computer controlled high resolution video and confocal laser scanning microscopy, we examined transfected samples for subcellular localization of the chimeric GFP-GR protein. We observed significant fluorescence in the cytoplasm of about ten percent of total cells, approximately the fraction that  
30 typically acquires transfected DNA. Thus, the GFP was functional as a chromophore in a majority of the expressing cells in this mammalian system.



Upon exposure to dexamethasone, translocation of GFP-GR occurs in 100% of fluorescing cells, with the rate of cytoplasm-to-nuclear translocation dependent on the concentration of hormone. At 10 nM, complete translocation was induced within 10 min at 37°C, with half maximal nuclear accumulation at 5 min; this rate is consistent with previous findings (Picard, D. & Yamamoto, K. R. (1987) *EMBO. J.* 6, 3333-3340). The rate of translocation is decreased with 1 nM dexamethasone (complete translocation over 30 min with half maximum at 9-10 min) and further reduced with 0.1 nM dexamethasone (complete translocation within 2 hours with half maximum at 1 hour).

Analysis of a time-lapse series revealed that GFP-GR accumulated along fibrillar structures and in the perinuclear region very rapidly after hormone addition, probably within seconds. Murine adenocarcinoma cells were cultured on cover slips and transfected with GFP-GR fusion chimera one day before microscopy. Cells were placed into a Dvorak-Stotler chamber and perfused at 37°C with assay buffer for 3 min, then with buffer containing 1 nM dexamethasone for 2 hrs. With real-time imaging, perinuclear accumulation was observed in a pulsatile pattern with 1-2 second intervals between brightness changes. GFP-GR accumulation was more intense along fibrillar structures in the perinuclear region. After 3 min with 1 nM dexamethasone, GFP-GR was noticeably present in the nucleus, but not in the nucleoli. When approximately 1/3 of the protein had been translocated (9-10 min), a punctate pattern appeared, and translocation was complete after 30 min. During translocation, the cells frequently became rounded, and moved along the long axis of the cell. We observed reduction of the cell surface, as well as the nuclear volume during the translocation. One hour after hormone treatment, the cells reattach and regain a more flattened shape.

25

**Ligand Specificity of Cytoplasm-to-Nuclear Translocation.** Cells were treated with buffer (A), 10 nM 17- $\beta$ -estradiol (B), 10 nM dexamethasone (C), or 10 nM RU486 (D) for 30 min at 37°C. At the end of hormone treatment, images from living cells expressing GFP-GR were visualized with confocal laser scanning microscope as described above. While complete translocation of GFP-GR was observed in all fluorescing cells treated with dexamethasone, other classes of steroid hormones induced

30

GFP-GR translocation to varying extent reflective of the affinity for GR. The glucocorticoid antagonist RU486, known to have a high affinity for GR (Chakraborti, P. K., Garabedian, M. J., Yamamoto, K. R. & Simons, S. S. J. (1991) *J. Biol. Chem.* **266**, 22075-22078), was as potent as dexamethasone for induction of translocation.

- 5 Progesterone, a weak GR agonist, required a concentration 100-fold higher than dexamethasone for translocation; however, approximately 1/2 of the GFP-GR remained in the cytoplasm. In contrast, 17- $\beta$ -estradiol, a steroid hormone that does not bind GR, did not cause intranuclear GFP-GR accumulation (10 nM). Thus, GFP-GR maintained ligand-dependent cytoplasm-to-nuclear translocation, with analog specificity identical to
- 10 that for the untagged GR with the C656G point mutation (Chakraborti, P. K., Garabedian, M. J., Yamamoto, K. R. & Simons, S. S. J. (1991) *J. Biol. Chem.* **266**, 22075-22078). Furthermore, while ligand binding may suffice to trigger efficient cytoplasm-to-nuclear translocation, it may not necessarily cause binding of the receptor to its nuclear target and it may cause varying degrees of activation of the target gene.

15

- Role of Intracellular Free Calcium and Energy in GFP-GR Translocation.** Two important issues concerning the nuclear import of proteins were also addressed; these include the role of  $\text{Ca}^{++}$ , and the energy requirement of translocation. Intracellular stores of  $\text{Ca}^{++}$  were depleted by incubating the cells for 1 hour with the endoplasmic
- 20 reticulum  $\text{Ca}^{++}$ -ATPase inhibitor, thapsigargin, and the calcium ionophore, ionomycin, in calcium-free media (intracellular free calcium content was measured with ratio imaging in Fura-2 loaded cells). The cytoplasmic pattern of GFP-GR was not significantly altered by calcium depletion. When  $\text{Ca}^{++}$ -depleted cells were subsequently exposed to dexamethasone (10nM for 30 min. at 37°C) in  $\text{Ca}^{++}$ -free media, the hormone induced
- 25 complete translocation of GFP-GR, as seen by images taken from living cells with a confocal laser scanning microscope.

- To study the energy dependence of ligand binding, cells were exposed to dexamethasone (10nM) at 4°C; then hormone was removed and the cells were warmed
- 30 to 37°C under continuous monitoring with video-microscopy (cooled CCD camera from Axiovert 10 microscope system). At 4°C, translocation was completely arrested.

Rewarming led to complete translocation and reappearance of the focal GFP-GR localization. This experiment indicates that hormone binding to GR in living cells does not require energy in contrast to the energy-dependent step of translocation.

- 5 **Focal Accumulation of Nuclear GFP-GR Correlates with Transcriptional Activation.** When the intranuclear accumulation of GFP-GR is examined in detail, it is readily apparent that the receptor localizes most prominently at specific foci within the nucleus. In addition, there is a low level of accumulation in a diffuse reticular pattern, forming the basis for the nuclear background fluorescence. The number of these focal
- 10 accumulations are unique to dexamethasone-treated cells and are not observed in 17- $\beta$ -estradiol- or progesterone-exposed cells. In RU486-treated cells, focal points are not readily discernible. Instead, GFP-GR accumulates in a diffuse pattern with regions of condensation in a reticular pattern, such that regions of bright fluorescence appear thread-like in shape instead of as distinct foci. Depleting intracellular  $\text{Ca}^{++}$  did not affect
- 15 the dexamethasone-mediated formation of intranuclear foci. The ability of agonist to induce focal accumulation of GFP-GR correlated strongly with its ability to activate transcription (Fig. 3C). The striking accumulation of dexamethasone-activated GFP-GR into intranuclear foci immediately suggests that a specific architecture may underlie this distribution. To further examine the structure of intranuclear GR binding sites, confocal
- 20 laser scanning fluorescent microscopy and three dimensional image reconstruction was carried out.

#### **Organized Architecture of Interphase Nuclei As Revealed by GFP-GR.**

- Three-dimensional architecture of GFP-GR nuclear target sites was analyzed. Serial 0.5
- 25  $\mu\text{m}$  sections of nuclei from dexamethasone treated cells were collected with confocal laser scanning fluorescent microscope, digitized images were imported into a Silicon Graphics Indigo 2 workstation, and three dimensional image segmentation, rendering, analysis and reconstruction was carried out with the ANALYZE software. GFP-GR distributions in the nuclei are displayed as pseudocolored, voxel-gradient-shaded, three
- 30 dimensional projections. Three dimensional image analysis of the points of GR accumulation in dexamethasone-treated cells reveals a non-random distribution of

GFP-GR accumulation. Most strikingly, comparison of adjacent cells demonstrates a reproducible pattern of intranuclear structure for GFP-GR accumulation. A predominance of GFP-GR-accumulating foci is observed in the quadrant of the nucleus adjacent to the glass attachment surface of the cell, while a group of large patches of GFP-GR-containing foci are observed in the top half. Nucleolar structures were always devoid of GFP-GR. The nuclear pattern of RU486-treated cells was again strikingly different from dexamethasone-treated cells. Although essentially all of the GFP-GR is translocated, intranuclear RU486-liganded receptor is distributed throughout the nucleus in a reticular pattern but excluding nucleoli.

10

### 3134 cell line and derivatives

The 3134 cell line was derived from a mouse line designated 904.1. This cell was established by transfection of a murine mammary carcinoma line (C127) with a plasmid containing three functional segments: a) the bovine papilloma virus (BPV) 69% transforming fragment serves as a replicon in mammalian cells, b) the mouse mammary tumor virus (MMTV) LTR is a steroid responsive promoter and contains the GR binding sites, and c) the Ha-v-ras gene is a transforming oncogene and serves as a reporter for the MMTV promoter.

20

This plasmid replicates in 904.1 cells as an 9 kb episomal circle. During passage of this cell line, a spontaneous integration event occurred. This event resulted in the integration of a tandem array of the BPV/MMTV-LTR/Ha-v-ras cassette in perfect head-to-tail orientation. The integrated structure is diagrammed in Figure 2. Standard agarose gel electrophoresis and southern transfer hybridization analysis with a BPV probe (standard gel) was performed. CHEF gel high molecular weight analysis, again with a BPV probe (CHEF gel) was also performed. When the integrated array is digested with a one-cut restriction enzyme, the repeat unit of 8.8 kb is liberated as a single fragment (BamHI). NdeI and EcoRV (both no-cutters for the repeat unit) digestion leads to no digestion in the standard gel; EcoRV liberates a fragment approximately 2.2 megabase pairs in size. Enzymes that cut multiple times (PstI) give rise to the appropriate fragments for the repeat unit. Since the one cutter enzyme

30

releases only one size fragment from the array, the units must be organized in a perfect head-to-tail array.

Derivatives of 3134 cell lines containing pTET-nGFP-C656G and  
5 pOP-nGFP-C656G have been generated allowing GFP-GR expression upon tetracycline withdrawal from 5 ug/ml or upon induction with IPTG using standard procedures and as recommended by the manufacturer of these inducible systems, respectively. For example, 3616 is a single cell clone of 3134 with the pTET-nGFP-C656G DNA allowing acceptable level of GFP-GR expression that contains 200 copies of the MMTV  
10 LTR-cat-BPV tandem repeats; 3617 is another single cell clone of 3134 with the pTET-nGFP-C656G DNA but where the copy number of the MMTV LTR-cat-BPV tandem repeats dropped from 200 copies to 150 copies upon passage and then increased to about 170 copies in all cells upon further passage; 3596 is a single-cell clone of 3134 having 200 copies of the MMTV LTR-cat-BPV repeats and contains  
15 pOP-nGFP-C656G DNA; 3597 is a single cell clone of 3134 with 200 copies of the MMTV LTR-cat-BPV repeats and contains pOP-nGFP-C656G DNA. Acceptable levels of GFP-GR is reached in these derivatives after overnight induction of the regulatable promoters, e.g., after withdrawal of tetracycline in the case of 3616 and 3617, and after addition of IPTG in the case of 3596 and 3597.

20

#### **Gene Targeting with GFP-GR in 3134 Cell**

When the GFP-Glucocorticoid Receptor is transfected into the 3134 cell line, it is initially localized completely in the cytoplasm of the cell, as is normal, non-derivatized glucocorticoid receptor. When GFP-GR is activated with the GR ligand  
25 dexamethasone, the receptor translocates to the nucleus and accumulates on the BVP/MMTV-LTR/ras tandem array.

Figure 2 is a schematic representation of the appearance of GFP-GR when the fluorescent tag is excited by 489 nm laser light, and the 511 nm emission examined by  
30 confocal microscopy, using a standard fluorescein filter set. Individual sections are depicted for a representative 3134 nucleus. As one focuses on 0.18 micrometer sections

through the nucleus, a continuous fiber of intense light emission is detected over 5-7 sections, which corresponds to GFP-GR binding to the continuous BVP/MMTV-LTR/ras array.

## 5 Expression of GFP-GR in 3134 Cell

### 1) Materials.

#### a) Recombinant DNA

The LacSwitch<sup>TM</sup> inducible mammalian expression system (Stratagene catalog number 217450) is supplied with the phagemid DNAs: p3'SS for constitutive  
10 expression of the E. coli lac repressor and hygromycin resistance drug selectable marker; pOPI3 CAT for lac repressor regulated expression from the Rous sarcoma virus (RSV)-LTR promoter and neomycin resistance drug selectable marker; and pOPRSVI  
CAT for lac repressor regulated expression from the Rous sarcoma virus (RSV)-LTR promoter and neomycin resistance drug selectable marker.

15

Using standard recombinant DNA methodology, cDNA encoding GFP-GR fusion (either with the C656G mutation in the steroid binding domain of rat GR as present in the plasmid pCI-nGFP-C656G or wild type ligand binding domain of rat GR as present in the plasmid pCI-nGFP-rGR) is subcloned into the vector pOPRSVI CAT  
20 phagemid (Stratagene) at the NotI site. The cDNA should contain at its 5' end Kozak consensus sequence for efficient utilization of the translational initiation codon ATG, and at its 3' end the translational termination sequence followed by multiple translation termination sequences in all three reading frames. The vector provides RSV-LTR that has been engineered to be regulated by E. coli lac repressor so that in the mammalian  
25 cells with lac repressor, the strong RSV LTR promoter is inducible by the addition of 5 mM isopropyl-beta-D-thiogalactopyranoside (IPTG). The vector also provides an intron in the 5' untranslated region to ensure proper processing of the nascent transcript and maturation into productive mRNA. The vector also provides at the 3' untranslated region, herpes simplex thymidine kinase (TK) polyadenylation signal to ensure  
30 polyadenylation and message stability.

All DNAs used for introduction into mammalian cells are prepared from *E. coli* by alkali lysis procedure followed by isopycnic centrifugation by banding twice in cesium chloride/ethidium bromide gradient. Ethidium is removed by repeated extraction with isopentyl alcohol and cesium chloride by dialysis against large volumes of 10 mM Tris-HCl, pH 8/1mM EDTA (TE). After dialysis, DNA is extracted twice in phenol chloroform solution and then with chloroform before precipitation with 0.2 M sodium acetate (pH 5.5) and 70% ethanol at -20C. After collecting the precipitate by centrifugation, the DNA is washed with cold 70% ethanol, air dried, and then resuspended at a concentration of 1-2.5 mg/ml in TE.

10

#### b) Cell Line

The 3134 cell line contains a 9 kb repeat of the MMTV LTR fused to Ha-v-ras protooncogene in a BPV-based mammalian vector transformed into mouse carcinoma cell line C127.

15

Derivatives of 3134 cell lines containing pTET-nGFP-C656G and pOP-nGFP-C656G have been generated allowing GFP-GR expression upon tetracycline withdrawal from 5 ug/ml or upon induction with IPTG (about 1mM) using standard procedures and as recommended by the manufacturer of these inducible systems.

20 Acceptable levels of GFP-GR is reached in these derivative cells after overnight induction of the regulatable promoters.

#### 2) Maintenance of 3134 Cells

The 3134 cells are maintained in complete DMEM media [Dulbecco's Modified Eagle Medium (DMEM; Gibco-BRL catalog number 11965-084) supplement with 2 mM L-glutamine (Gibco-BRL catalog number 25030-024), 50ug/ml gentamicin reagent (Gibco-BRL catalog number 15750-011), and 10% fetal bovine serum (Gibco-BRL catalog number 26140-079) as monolayer in 162 cm<sup>2</sup> cell culture flasks (Costar catalog number 3150) at 37C in 5% CO<sub>2</sub> humidified air incubator. Upon confluence, cells are washed with Dulbecco's phosphate buffered saline (D-PBS) without calcium or magnesium and then treated with 6 ml of 0.05% trypsin/0.53 mM EDTA for several

30

minutes at RT (Gibco-BRL catalog number 25300-062). After cells round up, the side of the flask is tapped to dislodge cells from the flask surface. Repeated pipetting of the cells about three times results in a single cell suspension. One fifth of the cell suspension is transferred to a fresh 162 cm<sup>2</sup> flask containing 25 ml of the complete DMEM media.

- 5 The freshly diluted cells are returned to the 37C/5% CO<sub>2</sub> incubator.

### 3) Introduction of DNA into 3134 Cells by Electroporation

- 3134 cells grown to about 90% confluence are harvested using the trypsin-EDTA and the activity of the trypsin quenched by placing the single cell suspension into
- 10 at least equal volume of complete DMEM. Cells are counted in a hemacytometer and concentrated by centrifugation in bench top centrifuge (Sorvall RT6000D at 2,000 rpm for 5 minutes at 4C) after placing in 50 ml sterile conical tube (Falcon 2070). Supernatant is aspirated and cells are washed again using complete DMEM. After the second wash, the resulting cell pellet is resuspended at a concentration of 20 million
- 15 cells per 200 ul in cold DMEM and placed on ice. In 1.7 ml sterile microtube, appropriate DNA to be transfected, from 1-50ug of about 8 kb plasmid DNA (typically 1-20 ug), is placed. 200 ul 3134 cells is added and mixed well. The mixture is transferred to disposable electro chamber cuvettes (Gibco-BRL catalog number 11601-010) on ice. Cuvettes are placed into electroporation chamber and
- 20 electroporated (Gibco-BRL Cell-Porator catalog series 1600; settings: 250 volts, 800 or 1180 microFarads, low resistance). Recovery on ice is allowed for 5 minutes. Then, cells are removed from the cuvette and transferred to 50 ml conical tube with complete DMEM at RT. Cells are plated out on 150 x 25 mm tissue culture dish (Falcon 3025) in 30 ml of complete DMEM containing about one fourth the content of the
- 25 electroporation cuvette. Plates are placed in 37C/5% CO<sub>2</sub> incubator. Next day, media is changed to fresh media and selected with appropriate drug. Every two days media is changed and selection maintained with the appropriate drug.

### 4) Stable 3134 Cell Line Containing Lac Repressor

- 30 Electroporation of 1 ug, 5 ug, 10 ug, 30 ug, and 60 ug p3'SS DNA into 20 million 3134 cells in 200 ul cold DMEM is performed as described above. One day



following electroporation, cells are selected with 450 ug/ml hygromycin B (Calbiochem catalog number 40005) for about 10 days during which hygromycin resistant colonies arise. The colonies are isolated by treating with trypsin impregnated sterile 3 mm cloning discs (PGC Scientifics catalog number 09-060) and transferred into 24 well  
5 tissue culture cluster dish (Costar 3524; each well 16 mm) containing 2 ml of complete DMEM with 450ug/ml hygromycin B. The cells are allowed to grow in the dish in a 37C/5% CO<sub>2</sub> incubator until at least 50% confluence. Media is replaced every couple of days.

10 Cells are trypsinized and transferred to 100 x 20 mm dish (Falcon 3003) containing 10 ml of complete DMEM with 450 ug/ml hygromycin B. Then, at about confluence, cell are trypsinized and transferred to 150 x 25 mm dish with 30 ml of complete DMEM with 450 ug/ml hygromycin. Near confluence, cells were trypsinized and aliquot replated into 100 mm dish while the remaining cells are frozen down in 10%  
15 DMSO with complete DMEM at -70C. Cells grown on 100 mm dishes are harvested at confluence. Extracts are prepared and tested by Western blotting with rabbit anti-lac repressor polyclonal antibody (Stratagene catalog number 217449) and detected using goat anti-rabbit IgG horseradish peroxidase conjugated antibody (Bio-rad catalog number 170-6515) and enhanced chemiluminescent ECL western blotting detection  
20 reagents (Amersham catalog number RPN 2106). Cells expressing high amounts of the lac repressor are recloned and used for electroporating the pOPRSVI CAT plasmid containing GFP-C656G or GFP-rGR fusion cDNA.

5) Stable 3134 Cell Line Containing IPTG-Inducible GFP-GR or GFP-Fusion  
25 Protein Expression System.

3134 cells showing high levels of constitutive lac repressor expression without cytotoxic effects are used for electroporating pOPRSVI CAT plasmid containing the appropriate GFP-fusion protein cDNA. Following similar regiment as described above,  
30 the cells are selected with 450 ug/ml hygromycin B as well as 500 ug/ml geneticin (Gibco-BRL catalog number 11811-031). Following about ten days after selection,

again drug resistant colonies are isolated, expanded, and characterized. Characterization is achieved by examining uninduced as well as 5 mM IPTG induced cells using Southern blot analysis of the genomic DNA, Western blot analysis using antibody directed against GFP and GR or other steroid receptor, characterize for localization of the GFP-fusion protein on the MMTV LTR-tandem array upon exposure to the appropriate ligand. Cells with appropriate desirable characteristics are subcloned and serve as reagents for diagnostic purposes.

#### **Additional GFP-Steroid Fusion Proteins**

The same technology is used to prepare GFP fusions for each member of the steroid/thyroid receptor family, including the estrogen receptor, progesterone receptor, androgen receptor, mineralocorticoid receptor, thyroid receptor, retinoic acid receptor (RAR), RXR receptor, vitamin D receptor, and TCCD (dioxin) receptor. Chimeric proteins for each of the receptors is be prepared using GFP fused either to the N-terminus or the C-terminus of the appropriate receptor. The chimeras are tested for functional transcriptional transactivation activity, and for binding to cognate response elements using cell lines described below.

**GFP-ER:** For example, the glucocorticoid receptor portion of pCI-nGL1-C656G DNA has been replaced with a human estrogen receptor. The GFP-ER was found to be functional in both transcriptional activation as well as proper subcellular localization. This further demonstrates the utility of the expression vector as well as the GFP-tagging approach.

#### **Modification of the GFP fluorescent moiety**

The success of GFP-GR fusion in detecting and characterizing *in vivo* gene targeting indicates that modification and enhancement of the GFP polypeptide can be of considerable usefulness. These modifications include:

- 1) Changes in the excitation wavelength to permit activation of selected GFP fusions;
- 2) Changes in the emission wavelength to permit observation of selected GFP fusions (GFP's will emit with different colors);

3) Enhancements to the efficiency of excitation and emission (GFP's will be "brighter");

4) Enhancements of GFP-chimeric proteins production by engineering the GFP sequence for more efficient expression:

- 5 a) increasing the steady state level of GFP-chimeric protein mRNA levels through enhanced transcription, transcript processing, and RNA stability;
  - b) increasing translational efficiency of the GFP-chimeric proteins by conservative substitutions of the protein coding regions and/or by introducing codons preferred for translation in humans cells;
  - c) increasing stability of the GFP-chimeric proteins by identification of protein
  - 10 destabilizing sequences and selective elimination of such turnover signals;
- 5) Changes in GFP to define the minimal fluorescing polypeptide that could function in a fusion protein.

For example, in plasmids pCI-nGL2-C656G; pCI-nGL3-C656G; pCI-nGL4-C656G; pCI-nGL5-C656G; pCI-nGL7-C656G; pCI-nGL9-C656G; pCI-nGL10-C656G; pCI-nGL11-C656G, blue variants of GFP are used. Additionally, in these plasmids, humanized codons encoding the fusion protein were utilized.

#### Additional Cell Lines

20 Human and other mammalian cell lines can be prepared containing receptor binding sites (receptor response elements) in multimerized arrays for direct visualization of *in vivo* gene targeting.

- 1) For each receptor, these arrays can be created by gene amplification. The receptor response element is fused to the dihydrofolate reductase (dHFR) gene,
- 25 transfected into human cells, and integrants selected by resistance to methotrexate. After selection of the initial cell lines, further selection is applied using sequentially higher levels of methotrexate. This results in amplification of the dHFR/receptor response element sequence, providing an array to which the receptor/GFP fusion binding can be observed (directly analogous to the 3134 cell line for GR). Further details are
- 30 provided below.

2) An alternative approach is to multimerize the receptor response element by synthetic DNA synthesis, then introduce this amplified element directly into mammalian cells. This can provide a concentrated target for GFP-receptor localization.

## 5 **Generating Cells having Tandem Arrays**

An approach that is being used to generate tandem arrays in a re-engineered promoter to obtain a functional transcription unit:

- 1) the MMTV LTR in the plasmid pLTRLuc is being mutated by site directed mutagenesis to introduce appropriate restriction enzyme cleavage sites at approximately  
10 positions:
  - +110 from start of transcription (HindIII)
  - 40 from start of transcription (Sall)
  - 217 from start of transcription (XhoI)
  - 1100 from start of transcription (XmaI)
- 15 2) the MMTV LTR in the plasmid pLTRLuc is being mutated by site directed mutagenesis to introduce appropriate restriction enzyme cleavage sites at approximately positions:
  - +110 from start of transcription (HindIII)
  - 80 from start of transcription (Sall)
  - 20 -217 from start of transcription (XhoI)
  - 1100 from start of transcription (XmaI)
- 3) the DNA from (1) and (2) are being subcloned into two luciferase reporter gene vectors from Promega called pGL3-Basic (catalog number 88-1737) and pRL-CMV (catalog number 1068-2003) by using HindIII and XmaI digests of (1) and (2) and  
25 pGL3-Basic DNAs and inserting the mutated LTR fragments from (1) and (2) into the polylinker region of pGL3-Basic. In the case of pRL-CMV, this DNA is cleaved with PstI and BglII and the mutated LTR fragments from (1) and (2) inserted in place of the CMV Immediate Early Enhancer/Promoter regions in the pRL-CMV vector using a PstI/HindIII adapter on one end of the fragment. Also, before the insertion of the  
30 mutated LTR DNAs from (1) and (2), the Sall site present in the pGL3-Basic is

eliminated by digesting this vector with Sall, filling-in the end with DNA polymerase, and subsequent resealing of the filled-in end before transforming bacteria.

- 4) oligonucleotides containing the glucocorticoid response elements (GRE: 5' AGAACA<sub>nnn</sub>TGTTCT 3') (SEQ ID NO:4) or estrogen response elements (ERE: 5' AGGTC<sub>nnn</sub>TGACCT 3') (SEQ ID NO:5) are synthesized and then annealed (e.g., oligonucleotide 1 and 2; or oligonucleotide 3 and 4) such that one end contains a cohesive end for Sall and the other XhoI. Some examples of such oligonucleotides are (where the lower case letters are spacer bases and in the above designation would have been designated as "n"):

10 oligonucleotide 1: (SEQ ID NO:6):

5' tcgagcgcgcaAGAACA<sub>cag</sub>TGTTCTgacgacacgaAGAACA<sub>gga</sub>TGTTCTcgtacagtg 3'

oligonucleotide 2: (SEQ ID NO:7):

5' tcgacactgtacgAGAACA<sub>tcc</sub>TGTTCTtcgtgtcgtcAGAACA<sub>ctg</sub>TGTTCTtgcgcgc 3'

oligonucleotide 3: (SEQ ID NO:8):

15 5' tcgagcgcgcaAGGTC<sub>acag</sub>TGACCTgacgacacgaAGGTC<sub>agga</sub>TGACCTcgtacagtg 3'

oligonucleotide 4: (SEQ ID NO:9):

5' tcgacactgtacgAGGTC<sub>atcc</sub>TGACCTtcgtgtcgtcAGGTC<sub>actg</sub>TGACCTtgcgcgc 3'

- 5) the annealed oligonucleotides (1 and 2; or 3 and 4) are then ligated to generate large arrays in a perfect head-to-tail tandem array; these arrays may go through subcloning steps in bacteria to build larger arrays from smaller ones as well as to verify the integrity of the DNA sequence in the array by DNA sequencing.

6) the multimerized arrays from (5) are then inserted into the Sall/XhoI site in the mutated LTR of (3).

- 7) also, an additional series of constructs are also made lacking the region from -217 to

-1100 of the LTR by restriction enzyme digestion with Xho I and MluI for the pGL3-Basic vector derived clones of (6), filling in the XhoI/MluI cohesive ends, and resealing the blunt ends to generate a multimerized tandem array of a DNA binding site driving the expression of the luciferase gene only through the binding sites in the array and the region of the LTR containing the signal for the start point of transcription (the TATA box and the initiator sequence).

- 8) stable mammalian cell lines (e.g., from human origin or Chinese hamster ovary (CHO)) are generated with the DNAs from (6) and (7) after linearization of the DNA, e.g. with BamHI and KpnI for the pGL3-Basic based DNAs and BamHI for the pRL-CMV based DNAs, and transfecting or electroporating these DNAs into cells, as is commonly done to introduce foreign DNA into cell. Also included in the transfection or electroporation is a DNA for a drug selectable marker such as neomycin gene or hygromycin that allows for drug selection (e.g., G418 or hygromycin, respectively) of cells that have taken up the foreign DNA. In addition, green fluorescent protein (GFP)-tagged glucocorticoid, estrogen, and orphan receptors or DNA binding proteins appropriate for the factor DNA-binding DNA element is also co-transfected or electroporated.
- 9) isolated clones from drug selected cells from (8) are characterized for the number of copies and dispersal of the introduced DNAs from (6) and (7). Also, characterized is the amount of fluorescence derived from the GFP tagged-receptors or DNA-binding protein as well as the functionality of the expressed protein.
- 10) because most of the cell lines from (9) will have the foreign DNA integrated throughout the genome, localization of the GFP-tagged DNA binding molecule on the arrays in each of these transcription unit would result in intense fluorescence spot or focus. It is also possible that an extremely few number of cells will have integrated the foreign DNA in small arrays which should look like slight elongated intense spot. The desirable cells are those that contain visible intense foci due to binding of the GFP-tagged DNA binding protein on the transcription unit. In the case of the glucocorticoid receptor, the desirable cell lines are such that agonist dexamethasone treatment results in accumulation of the GFP-GR on the transcription unit to generate intense foci but not antagonist RU486 treatment.
- 11) DNAs from (6) or (7) will also be ligated to generate tandem arrays of each transcription unit and cloned into cosmid vector or phage P1 vector.
- 12) the tandem array transcription units will be treated as in (8).
- 13) the result of (11) will be characterized as in (9) and (10) to obtain cell lines with desirable properties.

14) DNA from (6) and (7) will be linked to a constitutively expressible dihydrofolate reductase (DHFR) gene by standard DNA subcloning techniques.

15) the DNA from (14) will be introduced into mammalian cells lacking the tumor suppressor gene, p53, using a similar procedure as in (8) except leaving out the drug  
5 selectable marker DNAs (due to the fact that DHFR gene is a drug selectable marker gene) and then selected for two weeks with methotrexate at a concentration about four times above the LD90. Selection media is replaced every 2-3 days. After methotrexate resistant colonies appear (in approximately two weeks), the colonies are pooled and part of the pool is frozen down while the remainder is used in a repeat selection with a  
10 concentration of methotrexate about four times higher than that previously used. This last selection procedure is repeated a number of times to finally select out cells with a highly amplified tandem array copies of DNAs containing the multimerized binding sites, the reporter gene, and the DHFR gene.

16) the cell lines after the last methotrexate selection are characterized for the  
15 presence of at least one tandem array and localization of the GFP-tagged DNA binding protein on this array analyzed. Localization on this array is manifested as a long linear highly fluorescent structure observed in the nucleus. In the case of GFP-GR, the localization on the array should occur upon treatment with agonist dexamethasone but not antagonist RU486.

20 17) other approaches to generating array envisioned is the use of site specific recombinases in vivo to generate precisely tailored tandem arrays.

18) while items 1-16 focuses on the use of a genetically engineered artificial promoter/enhancer to generate arrays, tandem arrays may be made from just binding sites only or from natural regulatory DNAs harboring desirable factor-DNA binding  
25 sites.

19) since the higher eukaryotic genomes contain naturally occurring repetitive sequences, interactions of appropriately tagged DNA-binding protein with such naturally occurring sequences may also prove useful.

Using this general approach, functional receptor/cell line pairs can be established for each receptor-GFP fusion. These reagents provide a simple, rapid, straightforward, sensitive, and biologically relevant assay for each receptor.

- 5           Using standard recombinant DNA methodology, DNA encoding GFP-fusion protein sequences can be placed under the control of high expression eukaryotic promoter/enhancers (e.g. CMV promoter/enhancer, SV40 promoter/enhancer, RSV LTR, herpes simplex thymidine kinase [TK] promoter, etc.), naturally occurring inducible promoter/enhancers (e.g. metallothionine promoter/enhancer, MMTV LTR, 10 heatshock promoter/enhancer, etc.), or synthetic inducible promoter/enhancers (e.g. GAL4-VP16 inducible system, Stratagene's LacSwitch<sup>TM</sup> inducible mammalian expression system (catalog number 217450), Life Technologies' Tet regulated expression system (catalog number 10583-011), etc.). These DNAs are introduced into mammalian cells along with DNA expressing a selectable marker (neomycin, 15 hygromycin, zeocin, etc.) or screenable marker (e.g. fluorescence, foci formation, etc.) by standard protocol (e.g. calcium phosphate co-precipitation, electroporation, liposome-mediated transfection, viral infection, etc.). Following recovery of the cells to the introduction of DNA, approximately 1-3 days later, selectable agent is applied in the case of drug selectable marker to select for cells with stable integration of the selectable 20 marker.

- Upon continuous culturing of the cells under selection condition for an additional week during which clonal populations of cells will arise, the resulting cells are then characterized for the presence of the stably integrated DNA by Southern blot analysis 25 and PCR analysis, for the expression of the GFP-fusion protein by Western blot analysis, flow cytometry, and microscopic examination, for participation in ligand-dependent translocation/transactivation by assaying the activity of reporter genes (e.g. chloramphenicol acetyltransferase, luciferase, beta-galactosidase, etc.) under the control of the appropriate ligand inducible promoter/enhancer (e.g., glucocorticoid 30 response element containing promoter, estrogen response element containing promoter, etc.), and for localization on the target sequences in nuclei of expressing cells by high



resolution fluorescence imaging systems (e.g. confocal laser scanning microscopy, cooled CCD camera microscopy, etc.). Alternatively, stable transformants expressing GFP-fusion proteins can be directly isolated by fluorescence activated cell sorting (FACS) using appropriate excitation wavelengths and emission detector.

5

Although less preferable at present, an alternative to isolating clones of stable transformants will be the isolation of pools of stable transformants. The use of cell line/expression system combination may necessitate such a generation of a pool of stable transformants.

10

Additionally, while it is presently most efficient to obtain stable cell lines, GFP-fusion expression plasmid can be transiently introduced into cells and analyzed for ligand- dependent translocation/nuclear targeting.

#### 15 Use of Reagents

Because the use of GFP as a tag involves fusing a rather large protein (27 kDa) to GR, it was examined whether any GR activity has been compromised by GFP. To this end, GFP was fused to a rat GR with the C656G point mutation. This allowed selectively activation of GFP-GR without activating the endogenous receptor, and, thus, to assess the activity of the chimeric receptor independently of the endogenous GR. From the dose response curve and transcriptional activation of the MMTV LTR reporter gene (Fig. 3B), it is clear that GFP-GR can be selectively activated without activating the endogenous receptor.

25

By a number of criteria, GFP-GR functions very much like GR. In particular, the tagged receptor resides in the cytoplasm until activated by a ligand; it then translocates into the nucleus at a rate comparable to that previously reported (22). The rate and extent of GFP-GR translocation shows a dependence on the concentration of the activating ligand as well as a ligand specificity reflective of the native receptor.

30

Furthermore, since both dexamethasone and RU486 treatment lead to complete translocation of GFP-GR from the cytoplasm to the nucleus in all cells, essentially all of

the GFP-GR molecules exist in a conformation competent for both ligand binding and nuclear translocation. Once in the nucleus, GFP-GR's ability to activate the transcription of a MMTV LTR reporter gene depends on the type of activating ligand, consistent with previous results for GR. In the case of a potent agonist, dexamethasone, less GFP-GR is required for activation of transiently introduced MMTV LTRLuc reporter gene than for the endogenous GR, indicating that even with respect to transactivation potential, the presence of GFP has not altered the transcriptional potency ascribed to the C656G point mutation. Thus, in all aspects, the presence of GFP appears not to have affected normal GR function and has a general utility for studying the mechanisms of transcriptional regulation *in vivo*.

Since the S65T variant of GFP used here is highly excitable at 489 nm wavelength and resistant to photobleaching, it was possible to follow the course of cytoplasm-to-nuclear translocation of GR in a single living cell for an extended period of time. Upon binding to dexamethasone, GFP-GR moves vectorially toward the nucleus. Inhibition of import by chilling indicates that this transport is facilitated. Accumulation of GFP-GR along fibrillar structures before dexamethasone addition, and in the perinuclear region after dexamethasone addition, suggests that the cytoskeleton is involved in the transport process. Finally, the pulsatile brightness changes in the perinuclear region support an energy- and microtubule-dependent active transport process.

The rate of translocation of GFP-GR was dependent on hormone concentration, reflecting the dose- and time-dependence of GR action. This suggests that the rate of translocation contributes to GR function. The present invention indicates that cells having multiple copies of the response element in fluorescently detectable array can be a useful model to study reagents that modify rates of nuclear translocation and response element binding.

Finally, it is well-known that DNA in the nucleus is localized in a non-random fashion. The study of the pattern of nuclear fluorescence with GFP-GR supports the

idea of an inherent order in the organization of the interphase nuclei and may reflect structures related to this organization. The reproducibility in the pattern of GFP-GR accumulation between neighboring nuclei reflects an inherent order of the interphase nucleus with regards to both GR-target site architecture (the foci of bright fluorescence) as well as transcriptionally incompetent GFP-GR, which appears to accumulate in a reticular pattern, reminiscent of association with the nuclear matrix. Thus, functional differences in GFP-GR due to hormone-specific effects (e.g., dexamethasone vs. RU486) are reflected in the patterns of GFP-GR intranuclear accumulation, indicating a general utility of understanding intranuclear localization of GR in addressing hormone-mediated actions.

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

Although the present process has been described with reference to specific details of certain embodiments thereof, it is not intended that such details should be regarded as limitations upon the scope of the invention except as and to the extent that they are included in the accompanying claims.

## Sequence Listing

## (1) GENERAL INFORMATION:

(i) APPLICANT: The Government of the United States of America  
as represented by the Secretary  
Department of Health and Human Services  
Washington, D.C.

Htun Ph.D., Han  
Hager Ph.D., Gordon L.

(ii) TITLE OF INVENTION: METHODS AND COMPOSITIONS FOR MONITORING  
DNA BINDING MOLECULES IN LIVING CELLS

(iii) NUMBER OF SEQUENCES: 9

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Needle & Rosenberg  
(B) STREET: 127 Peachtree Street, Suite 1200  
(C) CITY: Atlanta  
(D) STATE: Georgia  
(E) COUNTRY: USA  
(F) ZIP: 30303

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: U.S. Serial No.60/008,373  
(B) FILING DATE: 08 Dec 1995  
(C) CLASSIFICATION:

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Selby, Elizabeth  
(B) REGISTRATION NUMBER: 38298  
(C) REFERENCE/DOCKET NUMBER: 14014.0183

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 404-688-0770  
(B) TELEFAX: 404-688-9880

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7257 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double

(ii) MOLECULE TYPE: nucleic acid

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1072..4284

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCAATATTGG CCATTAGCCA TATTATTCAT TGGTTATATA GCATAAATCA ATATTGGCTA	60
TTGGCCATTG CATACGTTGT ATCTATATCA TAATATGTAC ATTTATATTG GTCATGTCC	120
AATATGACCG CCATGTTGGC ATTGATTATT GACTAGTTAT TAATAGTAAT CAATTACGGG	180
GTCATTAGTT CATAGCCCAT ATATGGAGTT CCGCGTTACA TAACTTACGG TAAATGGCCC	240
GCCTGGCTGA CCGCCCAACG ACCCCCGCCC ATTGACGTCA ATAATGACGT ATGTTCCCAT	300
AGTAACGCCA ATAGGGACTT TCCATTGACG TCAATGGGTG GAGTATTTAC GGTAAACTGC	360
CCACTTGGCA GTACATCAAG TGTATCATAT GCCAAGTCCG CCCCTATTG ACGTCAATGA	420
CGGTAAATGG CCCGCTGGC ATTATGCCCA GTACATGACC TTACGGGACT TTCCTACTTG	480
GCAGTACATC TACGTATTAG TCATCGCTAT TACCATGGTG ATGCGGTTTT GGCAGTACAC	540
CAATGGGCGT GGATAGCGGT TTGACTCAGG GGGATTTCCA AGTCTCCACC CCATTGACGT	600
CAATGGGAGT TTGTTTTGGC ACCAAAATCA ACGGGACTTT CCAAATGTC GTAATAACCC	660
CGCCCCGTG ACGCAAATGG GCGGTAGGCG TGTACGGTGG GAGGTCTATA TAAGCAGAGC	720
TCGTTTAGTG AACCGTCAGA TACTAGAAG CTTTATTGCG GTAGTTTATC ACAGTTAAAT	780
TGCTAACGCA GTCAGTGCTT CTGACACAAC AGTCTCGAAC TTAAGCTGCA GAAGTTGGTC	840
GTGAGGCACT GGGCAGGTAA GTATCAAGGT TACAAGACAG GTTTAAGGAG ACCAATAGAA	900
ACTGGGCTTG TCGAGACAGA GAAGACTCTT GCGTTTCTGA TAGGCACCTA TTGGTCTTAC	960
TGACATCCAC TTTGCCTTTC TCTCCACAGG TGTCCACTCC CAGTTCAATT ACAGCTCTTA	1020
AGGCTAGAGT ACTTAATACG ACTCACTATA GGCTAGCGAA GGAGATCCGC C ATG GCC	1077
Met Ala	
1	
CAC CAT CAC CAC CAT CAC GGA TAT CCA TAC GAC GTG CCA GAT TAC GCT	1125
His His His His His His Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala	
5 10 15	
CAG TCG AGT GCC ATG AGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC	1173
Gln Ser Ser Ala Met Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val	
20 25 30	
CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTT TCT	1221
Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser	
35 40 45 50	
GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT	1269
Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu	
55 60 65	

68

AAA TTT ATT TGC ACT ACT GGA AAA CTA CCT GTT CCT TGG CCA ACA CTT	1317
Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu	
70 75 80	
GTC ACT ACT TTC ACT TAT GGT GTT CAA TGC TTT TCA AGA TAC CCA GAT	1365
Val Thr Thr Phe Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp	
85 90 95	
CAT ATG AAA CAG CAT GAC TTT TTC AAG AGT GCC ATG CCC GAA GGT TAT	1413
His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr	
100 105 110	
GTA CAG GAA AGA ACT ATA TTT TTC AAA GAT GAC GGG AAC TAC AAG ACA	1461
Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr	
115 120 125 130	
CGT GCT GAA GTC AAG TTT GAA GGT GAT ACC CTT GTT AAT AGA ATC GAG	1509
Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu	
135 140 145	
TTA AAA GGT ATT GAT TTT AAA GAA GAT GGA AAC ATT CTT GGA CAC AAA	1557
Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys	
150 155 160	
TTG GAA TAC AAC TAT AAC TCA CAC AAT GTA TAC ATC ATG GCA GAC AAA	1605
Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys	
165 170 175	
CAA AAG AAT GGA ATC AAA GTT AAC TTC AAA ATT AGA CAC AAC ATT GAA	1653
Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu	
180 185 190	
GAT GGA AGC GTT CAA CTA GCA GAC CAT TAT CAA CAA AAT ACT CCA ATT	1701
Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile	
195 200 205 210	
GGC GAT GGC CCT GTC CTT TTA CCA GAC AAC CAT TAC CTG TCC ACA CAA	1749
Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln	
215 220 225	
TCT GCC CTT TCG AAA GAT CCC AAC GAA AAG AGA GAC CAC ATG GTC CTT	1797
Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu	
230 235 240	
CTT GAG TTT GTA ACA GCT GCT GGG ATT ACA CAT GGC ATG GAT GAA CTA	1845
Leu Glu Phe Val Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu	
245 250 255	
TAC AAA GGC GCC GGC GCT GGT GCT GGT GCT GGC GCC ATC AGC GCG CTG	1893
Tyr Lys Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Ile Ser Ala Leu	
260 265 270	
ATC CTG GAC TCC AAA GAA TCC TTA GCT CCC CCT GGT AGA GAC GAA GTC	1941
Ile Leu Asp Ser Lys Glu Ser Leu Ala Pro Pro Gly Arg Asp Glu Val	
275 280 285 290	
CCT GGC AGT TTG CTT GGC CAG GGG AGG GGG AGC GTA ATG GAC TTT TAT	1989
Pro Gly Ser Leu Leu Gly Gln Gly Arg Gly Ser Val Met Asp Phe Tyr	
295 300 305	

AAA AGC CTG AGG GGA GGA GCT ACA GTC AAG GTT TCT GCA TCT TCG CCC Lys Ser Leu Arg Gly Gly Ala Thr Val Lys Val Ser Ala Ser Ser Pro 310 315 320	2037
TCA GTG GCT GCT GCT TCT CAG GCA GAT TCC AAG CAG CAG AGG ATT CTC Ser Val Ala Ala Ala Ser Gln Ala Asp Ser Lys Gln Gln Arg Ile Leu 325 330 335	2085
CTT GAT TTC TCG AAA GGC TCC ACA AGC AAT GTG CAG CAG CGA CAG CAG Leu Asp Phe Ser Lys Gly Ser Thr Ser Asn Val Gln Gln Arg Gln Gln 340 345 350	2133
CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln 355 360 365 370	2181
CAG CCA GGC TTA TCC AAA GCC GTT TCA CTG TCC ATG GGG CTG TAT ATG Gln Pro Gly Leu Ser Lys Ala Val Ser Leu Ser Met Gly Leu Tyr Met 375 380 385	2229
GGA GAG ACA GAA ACA AAA GTG ATG GGG AAT GAC TTG GGC TAC CCA CAG Gly Glu Thr Glu Thr Lys Val Met Gly Asn Asp Leu Gly Tyr Pro Gln 390 395 400	2277
CAG GGC CAA CTT GGC CTT TCC TCT GGG GAA ACA GAC TTT CGG CTT CTG Gln Gly Gln Leu Gly Leu Ser Gly Glu Thr Asp Phe Arg Leu Leu 405 410 415	2325
GAA GAA AGC ATT GCA AAC CTC AAT AGG TCG ACC AGC GTT CCA GAG AAC Glu Glu Ser Ile Ala Asn Leu Asn Arg Ser Thr Ser Val Pro Glu Asn 420 425 430	2373
CCC AAG AGT TCA ACG TCT GCA ACT GGG TGT GCT ACC CCG ACA GAG AAG Pro Lys Ser Ser Thr Ser Ala Thr Gly Cys Ala Thr Pro Thr Glu Lys 435 440 445 450	2421
GAG TTT CCC AAA ACT CAC TCG GAT GCA TCT TCA GAA CAG CAA AAT CGA Glu Phe Pro Lys Thr His Ser Asp Ala Ser Ser Glu Gln Gln Asn Arg 455 460 465	2469
AAA AGC CAG ACC GGC ACC AAC GGA GGC AGT GTG AAA TTG TAT CCC ACA Lys Ser Gln Thr Gly Thr Asn Gly Gly Ser Val Lys Leu Tyr Pro Thr 470 475 480	2517
GAC CAA AGC ACC TTT GAC CTC TTG AAG GAT TTG GAG TTT TCC GCT GGG Asp Gln Ser Thr Phe Asp Leu Leu Lys Asp Leu Glu Phe Ser Ala Gly 485 490 495	2565
TCC CCA AGT AAA GAC ACA AAC GAG AGT CCC TGG AGA TCA GAT CTG TTG Ser Pro Ser Lys Asp Thr Asn Glu Ser Pro Trp Arg Ser Asp Leu Leu 500 505 510	2613
ATA GAT GAA AAC TTG CTT TCT CCT TTG GCG GGA GAA GAT GAT CCA TTC Ile Asp Glu Asn Leu Leu Ser Pro Leu Ala Gly Glu Asp Asp Pro Phe 515 520 525 530	2661
CTT CTC GAA GGG AAC ACG AAT GAG GAT TGT AAG CCT CTT ATT TTA CCG Leu Leu Glu Gly Asn Thr Asn Glu Asp Cys Lys Pro Leu Ile Leu Pro 535 540 545	2709

70

GAC ACT AAA CCT AAA ATT AAG GAT ACT GGA GAT ACA ATC TTA TCA AGT Asp Thr Lys Pro Lys Ile Lys Asp Thr Gly Asp Thr Ile Leu Ser Ser 550 555 560	2757
CCC AGC AGT GTG GCA CTA CCC CAA GTG AAA ACA GAA AAA GAT GAT TTC Pro Ser Ser Val Ala Leu Pro Gln Val Lys Thr Glu Lys Asp Asp Phe 565 570 575	2805
ATT GAA CTT TGC ACC CCC GGG GTA ATT AAG CAA GAG AAA CTG GGC CCA Ile Glu Leu Cys Thr Pro Gly Val Ile Lys Gln Glu Lys Leu Gly Pro 580 585 590	2853
GTT TAT TGT CAG GCA AGC TTT TCT GGG ACA AAT ATA ATT GGT AAT AAA Val Tyr Cys Gln Ala Ser Phe Ser Gly Thr Asn Ile Ile Gly Asn Lys 595 600 605 610	2901
ATG TCT GCC ATT TCT GTT CAT GGT GTG AGT ACC TCT GGA GGA CAG ATG Met Ser Ala Ile Ser Val His Gly Val Ser Thr Ser Gly Gly Gln Met 615 620 625	2949
TAC CAC TAT GAC ATG AAT ACA GCA TCC CTT TCT CAG CAG CAG GAT CAG Tyr His Tyr Asp Met Asn Thr Ala Ser Leu Ser Gln Gln Gln Asp Gln 630 635 640	2997
AAG CCT GTT TTT AAT GTC ATT CCA CCA ATT CCT GTT GGT TCT GAA AAC Lys Pro Val Phe Asn Val Ile Pro Pro Ile Pro Val Gly Ser Glu Asn 645 650 655	3045
TGG AAT AGG TGC CAA GGC TCC GGA GAG GAC AGC CTG ACT TCC TTG GGG Trp Asn Arg Cys Gln Gly Ser Gly Glu Asp Ser Leu Thr Ser Leu Gly 660 665 670	3093
GCT CTG AAC TTC CCA GGC CGG TCA GTG TTT TCT AAT GGG TAC TCA AGC Ala Leu Asn Phe Pro Gly Arg Ser Val Phe Ser Asn Gly Tyr Ser Ser 675 680 685 690	3141
CCT GGA ATG AGA CCA GAT GTA AGC TCT CCT CCA TCC AGC TCG TCA GCA Pro Gly Met Arg Pro Asp Val Ser Ser Pro Pro Ser Ser Ser Ser Ala 695 700 705	3189
GCC ACG GGA CCA CCT CCC AAG CTC TGC CTG GTG TGC TCC GAT GAA GCT Ala Thr Gly Pro Pro Pro Lys Leu Cys Leu Val Cys Ser Asp Glu Ala 710 715 720	3237
TCA GGA TGT CAT TAC GGG GTG CTG ACA TGT GGA AGC TGC AAA GTA TTC Ser Gly Cys His Tyr Gly Val Leu Thr Cys Gly Ser Cys Lys Val Phe 725 730 735	3285
TTT AAA AGA GCA GTG GAA GGA CAG CAC AAT TAC CTT TGT GCT GGA AGA Phe Lys Arg Ala Val Glu Gly Gln His Asn Tyr Leu Cys Ala Gly Arg 740 745 750	3333
AAC GAT TGC ATC ATT GAT AAA ATT CGA AGG AAA AAC TGC CCA GCA TGC Asn Asp Cys Ile Ile Asp Lys Ile Arg Arg Lys Asn Cys Pro Ala Cys 755 760 765 770	3381
CGC TAT CGG AAA TGT CTT CAG GCT GGA ATG AAC CTT GAA GCT CGA AAA Arg Tyr Arg Lys Cys Leu Gln Ala Gly Met Asn Leu Glu Ala Arg Lys 775 780 785	3429
ACA AAG AAA AAA ATC AAA GGG ATT CAG CAA GCC ACT GCA GGA GTC TCA	3477





ACC TAC TGC TTC CAG ACA TTT TTG GAT AAG ACC ATG AGT ATT GAA TTC Thr Tyr Cys Phe Gln Thr Phe Leu Asp Lys Thr Met Ser Ile Glu Phe 1030 1035 1040	4197
CCA GAG ATG TTA GCT GAA ATC ATC ACT AAT CAG ATA CCA AAA TAT TCA Pro Glu Met Leu Ala Glu Ile Ile Thr Asn Gln Ile Pro Lys Tyr Ser 1045 1050 1055	4245
AAT GGA AAT ATC AAA AAG CTT CTG TTT CAT CAA AAA TGA CTGCCTTACT Asn Gly Asn Ile Lys Lys Leu Phe His Gln Lys * 1060 1065 1070	4294
AAGAAAGGTT GCCTTAAAGA AAGTTGAATT TATAGTCTAG AGTCGACCCG GCGGGCCGCT	4354
TCGAGCAGAC ATGATAAGAT ACATTGATGA GTTTGGACAA ACCACAAC TA GAATGCAGTG	4414
AAAAAAATGC TTTATTTGTG AAATTTGTGA TGCTATTGCT TTATTTGTAA CCATTATAAG	4474
CTGCAATAAA CAAGTTAACA ACAACAATTG CATTCAATTT ATGTTTCAGG TTCAGGGGGA	4534
GATGTGGGAG GTTTTTTAAA GCAAGTAAAA CCTCTACAAA TGTGGTAAAA TCGATAAGGA	4594
TCCGGGCTGG CGTAATAGCG AAGAGGCCCG CACCGATCGC CCTTCCCAAC AGTTGCGCAG	4654
CCTGAATGGC GAATGGACGC GCCCTGTAGC GGCGCATTAA GCGCGGCGGG TGTGGTGGTT	4714
ACGCGCAGCG TGACCGCTAC ACTTGCCAGC GCCCTAGCGC CCGCTCCTTT CGCTTTCTTC	4774
CCTTCCTTTC TCGCCACGTT CGCCGGCTTT CCCCCTCAAG CTCTAAATCG GGGGCTCCCT	4834
TTAGGGTTCC GATTTAGAGC TTACGGCAC CTCGACCGCA AAAA ACTTGA TTTGGGTGAT	4894
GGTTCACGTA GTGGGCCATC GCCCTGATAG ACGGTTTTTC GCCCTTTGAC GTTGGAGTCC	4954
ACGTTCTTTA ATAGTGGACT CTGTTCCAA ACTGGAACAA CACTCAACCC TATCTCGGTC	5014
TATCTTTTGG ATTTATAAGG GATTTTGCCG ATTTGCGCCT ATTGGTTAAA AAATGAGCTG	5074
ATTTAACAAA TATTTAACGC GAATTTTAAC AAAATATTAA CGTTTACAAT TTCGCCTGAT	5134
GCGGTATTTT CTCCTTACGC ATCTGTGCGG TATTTACAC ACATATGGT GCACTCTCAG	5194
TACAATCTGC TCTGATGCCG CATAGTTAAG CCAGCCCCGA CACCCGCCAA CACCCGCTGA	5254
CGCGCCCTGA CGGGCTTGTC TGCTCCCGGC ATCCGCTTAC AGACAAGCTG TGACCGTCTC	5314
CGGGAGCTGC ATGTGTCAGA GGTTTTCACC GTCATCACCG AAACGCGCGA GACGAAAGGG	5374
CCTCGTGATA CGCCTATTTT TATAGGTTAA TGTCATGATA ATAATGGTTT CTTAGACGTC	5434
AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTATTATTT TCTAAATACA	5494
TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT AATATTGAAA	5554
AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTCGCCCTT ATCCCTTTT TTGCGGCATT	5614
TTGCCCTCCT GTTTTGTCTC ACCCAGAAAC GCTGGTGAAA GTAAAAGATG CTGAAGATCA	5674
GTTGGGTGCA CGAGTGGGTT ACATCGAACT GGATCTCAAC AGCGGTAAGA TCCTTGAGAG	5734
TTTTCGCCCC GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCTGC TATGTGGCGC	5794

GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC ACTATTCTCA	5854
GAATGACTTG GTTGAGTACT CACCAGTCAC AGAAAAGCAT CTTACGGATG GCATGACAGT	5914
AAGAGAATTA TGCAAGTGTG CCATAACCAT GAGTGATAAC ACTGCGGCCA ACTTACTTCT	5974
GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTTTG CACAACATGG GGGATCATGT	6034
AACTCGCCTT GATCGTTGGG AACCGGAGCT GAATGAAGCC ATACCAAACG ACGAGCGTGA	6094
CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAAA CTATTAACTG GCGAACTACT	6154
TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG TTGCAGGACC	6214
ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG GAGCCGGTGA	6274
GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT CCCGTATCGT	6334
AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC AGATCGCTGA	6394
GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT CATATATACT	6454
TTAGATTGAT TTAAACTTC ATTTTAAAT TAAAAGGATC TAGGTGAAGA TCCTTTTGA	6514
TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTC CACTGAGCGT CAGACCCCGT	6574
AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT GCTGCTTGCA	6634
AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC TACCAACTCT	6694
TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC TTCTAGTGTA	6754
GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACCG CCTACATACC TCGCTCTGCT	6814
AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG GGTGGACTC	6874
AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGT CGTGACACA	6934
CCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG AGCTATGAGA	6994
AAGCGCCACG CTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG GCAGGGTCGG	7054
AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT ATAGTCCTGT	7114
CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG GGGGGCGGAG	7174
CCTATGGAAA AACGCCAGCA ACGCGGCCTT TTTACGGTTC CTGGCCTTTT GCTGGCCTTT	7234
TGCTCACATG GCTCGACAGA TCT	7257

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1071 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala His His His His His His Gly Tyr Pro Tyr Asp Val Pro Asp  
 1 5 10 15  
 Tyr Ala Gln Ser Ser Ala Met Ser Lys Gly Glu Glu Leu Phe Thr Gly  
 20 25 30  
 Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys  
 35 40 45  
 Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu  
 50 55 60  
 Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro  
 65 70 75 80  
 Thr Leu Val Thr Thr Phe Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr  
 85 90 95  
 Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu  
 100 105 110  
 Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr  
 115 120 125  
 Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg  
 130 135 140  
 Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly  
 145 150 155 160  
 His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala  
 165 170 175  
 Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn  
 180 185 190  
 Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr  
 195 200 205  
 Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser  
 210 215 220  
 Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met  
 225 230 235 240  
 Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr His Gly Met Asp  
 245 250 255  
 Glu Leu Tyr Lys Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Ile Ser  
 260 265 270  
 Ala Leu Ile Leu Asp Ser Lys Glu Ser Leu Ala Pro Pro Gly Arg Asp  
 275 280 285  
 Glu Val Pro Gly Ser Leu Leu Gly Gln Gly Arg Gly Ser Val Met Asp  
 290 295 300  
 Phe Tyr Lys Ser Leu Arg Gly Gly Ala Thr Val Lys Val Ser Ala Ser  
 305 310 315 320

75

Ser Pro Ser Val Ala Ala Ala Ser Gln Ala Asp Ser Lys Gln Gln Arg  
 325 330 335  
 Ile Leu Leu Asp Phe Ser Lys Gly Ser Thr Ser Asn Val Gln Gln Arg  
 340 345 350  
 Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln  
 355 360 365  
 Gln Gln Gln Pro Gly Leu Ser Lys Ala Val Ser Leu Ser Met Gly Leu  
 370 375 380  
 Tyr Met Gly Glu Thr Glu Thr Lys Val Met Gly Asn Asp Leu Gly Tyr  
 385 390 395 400  
 Pro Gln Gln Gly Gln Leu Gly Leu Ser Ser Gly Glu Thr Asp Phe Arg  
 405 410 415  
 Leu Leu Glu Glu Ser Ile Ala Asn Leu Asn Arg Ser Thr Ser Val Pro  
 420 425 430  
 Glu Asn Pro Lys Ser Ser Thr Ser Ala Thr Gly Cys Ala Thr Pro Thr  
 435 440 445  
 Glu Lys Glu Phe Pro Lys Thr His Ser Asp Ala Ser Ser Glu Gln Gln  
 450 455 460  
 Asn Arg Lys Ser Gln Thr Gly Thr Asn Gly Gly Ser Val Lys Leu Tyr  
 465 470 475 480  
 Pro Thr Asp Gln Ser Thr Phe Asp Leu Leu Lys Asp Leu Glu Phe Ser  
 485 490 495  
 Ala Gly Ser Pro Ser Lys Asp Thr Asn Glu Ser Pro Trp Arg Ser Asp  
 500 505 510  
 Leu Leu Ile Asp Glu Asn Leu Leu Ser Pro Leu Ala Gly Glu Asp Asp  
 515 520 525  
 Pro Phe Leu Leu Glu Gly Asn Thr Asn Glu Asp Cys Lys Pro Leu Ile  
 530 535 540  
 Leu Pro Asp Thr Lys Pro Lys Ile Lys Asp Thr Gly Asp Thr Ile Leu  
 545 550 555 560  
 Ser Ser Pro Ser Ser Val Ala Leu Pro Gln Val Lys Thr Glu Lys Asp  
 565 570 575  
 Asp Phe Ile Glu Leu Cys Thr Pro Gly Val Ile Lys Gln Glu Lys Leu  
 580 585 590  
 Gly Pro Val Tyr Cys Gln Ala Ser Phe Ser Gly Thr Asn Ile Ile Gly  
 595 600 605  
 Asn Lys Met Ser Ala Ile Ser Val His Gly Val Ser Thr Ser Gly Gly  
 610 615 620  
 Gln Met Tyr His Tyr Asp Met Asn Thr Ala Ser Leu Ser Gln Gln Gln  
 625 630 635 640

76

Asp Gln Lys Pro Val Phe Asn Val Ile Pro Pro Ile Pro Val Gly Ser  
 645 650 655  
 Glu Asn Trp Asn Arg Cys Gln Gly Ser Gly Glu Asp Ser Leu Thr Ser  
 660 665 670  
 Leu Gly Ala Leu Asn Phe Pro Gly Arg Ser Val Phe Ser Asn Gly Tyr  
 675 680 685  
 Ser Ser Pro Gly Met Arg Pro Asp Val Ser Ser Pro Pro Ser Ser Ser  
 690 695 700  
 Ser Ala Ala Thr Gly Pro Pro Pro Lys Leu Cys Leu Val Cys Ser Asp  
 705 710 715 720  
 Glu Ala Ser Gly Cys His Tyr Gly Val Leu Thr Cys Gly Ser Cys Lys  
 725 730 735  
 Val Phe Phe Lys Arg Ala Val Glu Gly Gln His Asn Tyr Leu Cys Ala  
 740 745 750  
 Gly Arg Asn Asp Cys Ile Ile Asp Lys Ile Arg Arg Lys Asn Cys Pro  
 755 760 765  
 Ala Cys Arg Tyr Arg Lys Cys Leu Gln Ala Gly Met Asn Leu Glu Ala  
 770 775 780  
 Arg Lys Thr Lys Lys Lys Ile Lys Gly Ile Gln Gln Ala Thr Ala Gly  
 785 790 795 800  
 Val Ser Gln Asp Thr Ser Glu Asn Pro Asn Lys Thr Ile Val Pro Ala  
 805 810 815  
 Ala Leu Pro Gln Leu Thr Pro Thr Leu Val Ser Leu Leu Glu Val Ile  
 820 825 830  
 Glu Pro Glu Val Leu Tyr Ala Gly Tyr Asp Ser Ser Val Pro Asp Ser  
 835 840 845  
 Ala Trp Arg Ile Met Thr Thr Leu Asn Met Leu Gly Gly Arg Gln Val  
 850 855 860  
 Ile Ala Ala Val Lys Trp Ala Lys Ala Ile Leu Gly Leu Arg Asn Leu  
 865 870 875 880  
 His Leu Asp Asp Gln Met Thr Leu Leu Gln Tyr Ser Trp Met Phe Leu  
 885 890 895  
 Met Ala Phe Ala Leu Gly Trp Arg Ser Tyr Arg Gln Ser Ser Gly Asn  
 900 905 910  
 Leu Leu Cys Phe Ala Pro Asp Leu Ile Ile Asn Glu Gln Arg Met Ser  
 915 920 925  
 Leu Pro Gly Met Tyr Asp Gln Cys Lys His Met Leu Phe Val Ser Ser  
 930 935 940  
 Glu Leu Gln Arg Leu Gln Val Ser Tyr Glu Glu Tyr Leu Cys Met Lys  
 945 950 955 960

77

Thr Leu Leu Leu Leu Ser Ser Val Pro Lys Glu Gly Leu Lys Ser Gln  
                   965                                  970                                  975  
 Glu Leu Phe Asp Glu Ile Arg Met Thr Tyr Ile Lys Glu Leu Gly Lys  
                   980                                  985                                  990  
 Ala Ile Val Lys Arg Glu Gly Asn Ser Ser Gln Asn Trp Gln Arg Phe  
                   995                                  1000                                  1005  
 Tyr Gln Leu Thr Lys Leu Leu Asp Ser Met His Glu Val Val Glu Asn  
                   1010                                  1015                                  1020  
 Leu Leu Thr Tyr Cys Phe Gln Thr Phe Leu Asp Lys Thr Met Ser Ile  
                   1025                                  1030                                  1035                                  1040  
 Glu Phe Pro Glu Met Leu Ala Glu Ile Ile Thr Asn Gln Ile Pro Lys  
                   1045                                  1050                                  1055  
 Tyr Ser Asn Gly Asn Ile Lys Lys Leu Leu Phe His Gln Lys \*  
                   1060                                  1065                                  1070

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCGCGCTGAT CAGAATTCCT TTAGGAATT CTGATCAGCG CGCTGA

46

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AGAACANNNT GTTCT

15

## (2) INFORMATION FOR SEQ ID NO:5:

78

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AGGTCANNNT GACCT

15

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TCGAGCGCGC AAGAACACAG TGTTCGACG ACACGAAGAA CAGGATGTTT TCGTACAGTG

60

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TCGACACTGT ACGAGAACAT CCTGTTCTTC GTGTCGTCAG AACACTGTGT TCTTGCGCGC

60

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide



(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCGAGCGCGC AAGGTCACAG TGACCTGACG ACACGAAGGT CAGGATGACC TCGTACAGTG 60

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TCGACACTGT ACGAGGTCAT CCTGACCTTC GTGTCGTCAG GTCAGTGTGA CCTTGCGCGC 60

**What is claimed is:**

1. A mammalian cell having a plurality of steroid receptor response elements in an array such that the response element can be directly detected when bound by fluorescently labeled steroid receptor.
2. The cell of claim 1, wherein the response element is integrated into the genome of the cell.
3. The cell of claim 1, wherein the response element is present in the mouse mammary tumor virus long terminal repeat.
4. The cell of claim 3, wherein the cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).
5. The cell of claim 1, further comprising a nucleic acid encoding a chimeric protein wherein a fluorescent protein is fused to the steroid receptor.
6. The cell of claim 5, wherein the nucleic acid is integrated into the genome of the cell.
7. The cell of claim 5, wherein the fluorescent protein is green fluorescent protein.
8. The cell of claim 5, wherein the steroid receptor is glucocorticoid receptor.
9. The cell of claim 8, wherein the chimeric protein has the amino acid sequence set forth in SEQ ID NO: 2.
10. A method of screening for a compound that binds to a selected nucleic acid comprising:
  - a. contacting compound fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of the nucleic acid in an array such that the nucleic

- acid can be directly detected when bound by fluorescently labeled compound;  
and
- b. directly detecting the location of fluorescence within the cell,  
fluorescence aggregated at the site of the nucleic acid array indicating a compound that binds  
to the selected nucleic acid.
11. The method of claim 10, wherein the compound is fluorescently labeled with a green  
fluorescent protein.
12. The method of claim 10, wherein the selected nucleic acid is integrated into the  
genome of the cell.
13. A method of screening for a ligand that activates gene targeting of a steroid receptor  
in the nucleus of a mammalian cell comprising:  
a. contacting the ligand with the cell of claim 5; and  
b. directly detecting the location of fluorescence within the cell,  
fluorescence aggregated at the site of the steroid receptor response element array in the  
nucleus indicating a ligand that activates the gene targeting of a steroid receptor in the  
nucleus of a mammalian cell.
14. The method of claim 13, wherein the fluorescent protein is green fluorescent protein.
15. The method of Claim 13, wherein the cell is a cell of the cell line designated 3134  
deposited with American Type Culture Collection under accession number \_ CRL-  
11998 (ATCC).
16. A method of screening for a ligand that activates the translocation of a steroid  
receptor to the nucleus in a mammalian cell comprising:  
a. contacting the cell of claim 5 with the ligand; and  
b. directly detecting the location of fluorescence within the cell,

a change in the relative fluorescence of the nucleus to the cytoplasm such as to increase the fluorescence of the nucleus indicating a ligand that activates the translocation of a steroid receptor to the nucleus in a mammalian cell.

17. The method of claim 16, wherein the fluorescent protein is green fluorescent protein.
18. The method of claim 16, wherein the mammalian cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).
19. A method of detecting in a biological sample the presence of an agonist of a steroid receptor comprising:
  - a. contacting the sample with the cell of claim 5 and
  - b. directly detecting the location of fluorescence within the cell,the location of fluorescence aggregated at the site of the steroid receptor response element array in the nucleus indicating the presence of an agonist of the steroid receptor in the sample.
20. The method of claim 19, wherein the fluorescent protein is green fluorescent protein.
21. The method of claim 19, wherein the cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).
22. A method of detecting in a biological sample the presence of an antagonist of a steroid receptor comprising:
  - a. contacting the sample and an agonist of the steroid receptor with the cell of claim 5 ; and
  - b. directly detecting the location of fluorescence within the cell,

the absence of fluorescence substantially aggregated at the site of the steroid receptor response element array in the nucleus indicating the presence of an antagonist of the steroid receptor in the sample.

23. The method of claim 22, wherein the fluorescent protein is green fluorescent protein.

24. The method of claim 22, wherein the cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).

25. A method of monitoring the level of an agonist of a steroid receptor in a subject comprising:

- a. periodically obtaining a biological sample from the subject,
- b. contacting the sample with the cell of claim 5, and
- c. directly detecting the location of fluorescence within the cell,

a decrease in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating a decrease in level of the steroid agonist of the steroid receptor in the sample and an increase in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid agonist of the steroid receptor in the sample.

26. The method of claim 25, wherein the fluorescent protein is green fluorescent protein.

27. The method of claim 25, wherein the cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).

28. A method of monitoring the balance between levels of an agonist of a steroid receptor and an antagonist of the steroid receptor in a subject comprising:

- a. periodically obtaining a biological sample from the subject,

- b. contacting the sample with the cell of claim 5, and
- c. directly detecting the location of fluorescence within the cell,

an increase in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid agonist relative to level of the steroid antagonist in the sample, and a decrease in fluorescence aggregated at the site of the steroid receptor response element in the nucleus in a later-obtained sample relative to an earlier-obtained sample indicating an increase in level of the steroid antagonist of the steroid receptor relative to level of the steroid agonist in the sample.

29. A method of determining an effective dosage of a steroid receptor agonist in a subject comprising:
- a. transferring into a set of cells from the patient a nucleic acid encoding a chimeric protein comprising a fluorescent protein fused to a steroid receptor;
  - b. contacting the cells in the set with one of a selected range of dosages of the steroid agonist; and
  - c. directly detecting location of fluorescence in the set of cells,
- a dosage capable of locating fluorescence substantially in the nucleus indicating an effective dosage of steroid receptor agonist.

30. The method of claim 28, wherein the fluorescent protein is green fluorescent protein.

31. A method of determining an effective dosage of a steroid receptor agonist to maintain steroid receptor activation for a selected period of time in a subject comprising:
- a. administering to the subject a dosage of the steroid receptor agonist,
  - b. periodically obtaining a biological sample from the subject,
  - c. contacting the sample with the cell of claim 5, and
  - d. directly detecting the location of fluorescence within the cell,
- a dosage that maintains the location of fluorescence at the site of the steroid receptor response element array in the nucleus for the selected period of time indicating an effective dosage.

32. The method of claim 31, wherein the fluorescent protein is green fluorescent protein.
33. The method of claim 31, wherein the cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).
34. A method of determining an effective dosage of a steroid receptor antagonist to abrogate agonist activity for a selected period of time in a subject comprising:
- administering to the subject a dosage of the steroid receptor agonist,
  - periodically obtaining a biological sample from the subject,
  - contacting the sample with the cell of claim 5, and
  - directly detecting the location of fluorescence within the cell,
  - a dosage that prevents the location of fluorescence at the site of the steroid receptor response element array in the nucleus for the selected period of time indicating an effective dosage.
35. The method of claim 34, wherein the fluorescent protein is green fluorescent protein.
36. The method of claim 34, wherein the cell is a cell of the cell line designated 3134 deposited with American Type Culture Collection under accession number CRL-11998 (ATCC).
37. A method of detecting a defect in a response pathway of a steroid receptor in a subject comprising transferring into a cell from the subject a nucleic acid functionally encoding a chimeric protein comprising a fluorescent protein fused to the steroid receptor and detecting the location of fluorescence within the cell as compared to the location of fluorescence within a normal, control cell transfected with the nucleic acid,
- a difference in location of fluorescence within the cell of the subject as compared to location of fluorescence within the normal, control cell indicating a defect in the response pathway of the steroid receptor.


38. A method of determining whether a defect in a response pathway of a steroid receptor in a subject is in translocation of the steroid receptor to a cell nucleus, comprising transferring into a cell from the subject having the defect a nucleic acid functionally encoding a chimeric protein comprising a fluorescent protein fused to the steroid receptor and detecting the location of fluorescence within the cell, the location of fluorescence substantially in the cytoplasm of the cell indicating the defect is in translocation of the steroid receptor to the nucleus.
39. A method of characterizing a ligand's effect on cellular localization of a compound to which the ligand binds in a cell comprising:
- contacting the ligand with a cell having the compound fluorescently labeled by a fluorescent protein and
  - directly detecting the location of fluorescence within the cell,
- the location of fluorescence in the cell indicating the localization effect of the ligand on the compound.
40. A method of determining a binding site for a DNA-binding protein comprising:
- contacting the DNA-binding protein fluorescently labeled by a fluorescent protein with a cell having a plurality of copies of a nucleic acid having a putative binding site in an array such that the putative binding site can be directly visualized when bound by the fluorescently labeled DNA-binding protein, and
  - directly detecting the location of fluorescence within the cell, the presence of fluorescence aggregated at the putative binding site indicating a binding site to which the DNA-binding protein binds.
41. A chimeric protein comprising a fluorescent protein fused to a transcription factor.
42. The protein of claim 41, wherein the transcription factor is a steroid receptor.





43. The protein of claim 41, wherein the fluorescent protein is a green fluorescent protein.
44. The protein of claim 43, wherein the green fluorescent protein is *Aequorea* green fluorescent protein.
45. The protein of claim 43, wherein the green fluorescent protein has a substitution of threonine for serine at amino acid 65.
46. The protein of claim 43, wherein the green fluorescent protein is fused to the transcription factor via a peptide linker.
47. The protein of claim 46, wherein the peptide linker comprises about five glycine-alanine repeating units.
48. The protein of claim 42, wherein the steroid receptor is glucocorticoid receptor and a green fluorescent protein is fused to the amino-terminal end of the glucocorticoid receptor.
49. The protein of claim 48, wherein the glucocorticoid receptor has a substitution of serine for cysteine 656.
50. The protein of claim 49 having the amino acid sequence set forth in SEQ ID NO: 2.
51. An isolated nucleic acid encoding the protein of claim 41.
52. An isolated nucleic acid encoding the protein of claim 42.
53. An isolated nucleic acid encoding the protein of claim 43.
54. An isolated nucleic acid encoding the protein of claim 48.

- 55. An isolated nucleic acid encoding the protein of claim 50.
- 56. An isolated nucleic acid encoding the protein of claim 50 having the nucleotide sequence set forth in SEQ ID NO: 1.
- 57. A cell containing the nucleic acid of claim 51.
- 58. A cell containing the nucleic acid of claim 55.

1/5  
FIG. 1



2/5

**Confocal Sections****FIG. 2**

3/5

FIG. 3A




FIG. 3B





FIG. 3C



# Nuclear Localization of GFP-GR in MCF7 Cells





FIG.4

5/5

# Nuclear Localization of GFP-ER



MDA-MB-231

MCF7

FIG.5